

***Mycobacterium gordonae* infection in freshwater fish from lakes and ponds in a park at São Paulo city, Brazil**

Ana Maria Cristina Rebello Pinto da Fonseca Martins^{1*}, Luara Lucena Cassiano²,
Marcia Helena Braga Catroxo³, Regina Stroebel⁴, Ana Paula Ruggiero-Couceiro⁵,
Carlos Eduardo Campos⁶, Paulo César Caldas⁷, Jesus Ramos⁸, Cristina Dib⁹,
Eliana Roxo¹⁰, Marcio Hipolito¹¹

^{1,2,3,9,10,11}Instituto Biológico de São Paulo/APTA/SAA, São Paulo, SP, Brazil.

⁴Prefeitura Municipal de São Paulo, SP, Brazil.

⁵Autonomous Veterinarian, São Bernardo do Campo, SP, Brazil.

^{6,7,8}Fundação Oswaldo Cruz, Rio de Janeiro, RJ, Brazil.

Abstract— In recent year's fish farming has greatly increased in Brazil, favoring the development of diseases such as mycobacteriosis. This is a chronic progressive disease that affects temperate and tropical fish, both freshwater and marine. Mycobacteriosis can occur in several species of fish and amphibian. In addition, some species of *Mycobacterium* spp. can be transmitted to humans by occupational or recreational source. A total of 54 fishes from lakes from São Paulo city, were collected and examined for mycobacteriosis. Granulomas were visualized in 5 fishes via histopathology (H&E), and acid-alcohol resistant bacilli were visualized in 8 animals by electron microscopy and 8 were positives using the Fite -Faraco technique. In this study, we isolated acid-fast bacillus from one fish which were identified as *M. gordonae* by molecular methods: PCR and sequencing.

Keywords— mycobacteriosis, pathology, aquaculture, sanity, disease.

I. INTRODUCTION

In recent years fish farming has greatly increased in Brazil, favoring the development of diseases such as mycobacteriosis (ISHIKAWA et al., 2001; ROMANO et al., 2012). This is a chronic progressive disease that affects temperate and tropical fish, both freshwater and marine (JACOBS et al., 2009).

Mycobacteria spp. can cause serious and costly diseases in different vertebrates and invertebrates, such as humans (tuberculosis, leprosy, Buruli ulcer), livestock (bovine tuberculosis) and ectotherms (reptiles, amphibians and fish) (BIET et al., 2005; GRANGE & YATES, 1986; JACOBS et al., 2009; REAVILL & SCHMIDT, 2012; SHINNICK & GOOD, 1994; TORTOLI, 2003; TURENNE et al., 2007).

The first *Mycobacterium* spp. was identified in carp in 1897 (BATAILLON, DUBARD, TERRE, 1897). This was named as *Mycobacterium piscium* and was shown to be highly pathogenic to frogs and some endothermic animals. The main species identified in captive and wild fishes are *M. marinum*, *M. fortuitum* and *M. chelonae* (mainly in marine fish). New species have also been proposed, including *M. salmoniphilum*. Other organisms related to *M. ulcerans* and the *M. tuberculosis* complex have also been recently implicated (GAUTHIER & RHODES, 2009; JACOBS et al., 2009).

Transmission typically occurs by ingestion of contaminated food and water, but transovarian transmission can also occur in viviparous species (ASTROFSKY et al., 2000; GAUTHIER & RHODES, 2009; JACOBS et al., 2009).

Granulomas are mainly located in the spleen, liver and kidney during the initial stages of the disease, but can spread to any other organs, leading to terminal illness. At the beginning of the infection, macrophages are invaded and become epithelioid cells. Giant cells may or not be present (GAUTHIER & RHODES, 2009; JACOBS et al., 2009).

In fish, the severity of mycobacteriosis ranges from chronic infection, without major changes in tissues and few losses, to severe and acute infection, with high mortality, depending on the mycobacteria and fish species involved. Clinical signs include weight loss, apathy and lethargy, decreased fertility spine defects, exophthalmia, abnormal behavioral, changes in skin color, and ulcerative lesions in the skin, gills, fins and musculature. There may be enlargement of liver, spleen, kidney and nodular lesions in internal organs (ASTROFSKY et al., 2000; GAUTHIER & RHODES, 2009; JACOBS et al., 2009; ROMANO et al., 2012).

There are few reports of mycobacteriosis in fish and amphibian species in Brazil. Studies are needed to understand the occurrence and consequences of the disease in animals maintained in captivity in lakes, ponds and parks (FERREIRA et al., 2006; ISHIKAWA et al., 2001; LEITE et al., 1998; ROMANO et al., 2012).

The aim of this study was to study this disease of fishes in the lakes and ponds park in São Paulo city and from decorative lake, unfit for consumption.

II. MATERIAL AND METHOD

2.1 Experimental Design

A total of 54 fishes (41 carps, 9 tilapias, 2 curimbatas (*Prochilodus lineatus*), 2 pirapitingas (*Piaractus brachipomus*), both male and female, were randomly collected from lakes and ponds in Jardim da Luz, located in downtown area at São Paulo City, Brazil. Samples of spleen, hepatopancreas, kidney and gills were fixed in 10% neutral buffered formalin or frozen. The sampled fishes varied in length from 10 cm to 52cm. Macroscopically, 2 carps showed lesions suggestive of granulomas.

2.2 H.E. Technique

Serial sections were prepared from the fixed material: fragments embedded paraffin. 5µm sections were cut using a microtome and adhered to the glass slides and stained by hematoxylin-eosin.

2.3 Fite- Faraco Ziehl-Neelsen technique (Z-N) (we used Fite-Faraco staining protocol, since the classic staining protocol of Ziehl Neelsen may result in false negatives).

Serial sections were prepared from the fixed material: fragments embedded paraffin. 5µm sections were cut using a microtome and adhered to the glass slides. The sections will be de-paraffinize in a solution composed of two parts of xylol and one part of peanut oil (or almond oil) for 15 minutes. The sections are then washed in tap water to remove the remaining xylene / oil mixture. Filter on carbol fuchsin solution, DO NOT HEAT, for 20 mins. Wash in running tap water. Differentiation will be done by means of 10% sulfuric acid for 2 minutes. Wash well in running tap water, rinse distilled water. Counterstain in 0.25% methylene blue for 20 seconds. Wash and blot dry. DO NOT DEHYDRATE IN ALCOHOL. Clear in xylene. Repeat the blotting-xylene treatment until section is clear. Mount in a DPX type mountan (FITE ET AL., 1947).

2.4 Negative Contrasting

The samples were suspended in 0.1M phosphate buffer pH 7.0 and placed in contact with metal grids previously coated with collodion and carbon film drained with filter paper. They were negatively contrasted with ammonium molybdate to 2% and pH 5.0 and observed using a Philips EM 208 (BRENNER & HORNE, 1959; HAYAT & MILLER, 1990) TEM.

2.5 PCR

For mycobacterial isolation, approximately one gram of each clinical specimen was ground with sterile sand, decontaminated by the classical Petroff method and seeded in four tubes containing medium of Stonebrink and four tubes with Petraghani medium. Two tubes of each seeded medium were incubated at 37°C and the remaining tubes at room temperature. All tubes were observed weekly for checking the growth of the colonies (KANTOR, 1988). To PCR, the isolated colonies were resuspended in 1.5mL sterile ultrapure water. DNA extraction was performed by inactivation of the samples by boiling at 100°C for 5 minutes, after which they were subjected to freezing at -20°C for at least one hour (BEMER-MELCHIOR & DRUGEON, 1999).

Thawed samples were PCR amplified using TB11-TB12 primers, designed for identification of the *Mycobacterium genus* (TELENTI et al., 1993). These generated a final product of 439 bases pairs. DNA amplifications were held in thermal cycler, submitting samples to the initial treatment of 95°C for 5 minutes, followed by 45 cycles of three temperatures: denaturation at 94°C for 1 minute, annealing at 65°C for 1 minute and extension at 72°C for 1 minute. After the last cycle, was held a final extension at 72°C for 7 minutes after which the product remained at 4°C until its analysis by electrophoresis in horizontal vat. PCR products were observed in 1.5% agarose gel containing 0.01% ethidium bromide, viewed under UV light and photographed with the aid of molecular gel doc system. To sequencing, the almost sequences of the 16S rDNA gene were obtained as described by CAMPOS et al. (2012).

III. RESULTS AND DISCUSSION

Of the 54 fishes examined, 8 were positives when stained using the Fite -Faraco Z-N technique (Fig 1). In the H&E staining, 5 animals presented numerous granulomas (Fig 2 a and b) of numerous sizes, with caseous necrosis in the center, eosinophilic cells and surrounded by inflammatory cells and fibroblasts (1 animal in the spleen, 3 kidneys and pancreas, 1 in hepatopancreas).




FIG 1 - PHOTOMICROGRAPH OF HEPATOPANCREAS SHOWING NUMEROUS RED MYCOBACTERIA (SMALL POINTS) INTO A GRANULOMA USING STAINED BY FITE- FARACO ZIEHL NIELSEN TECHNIQUE. (Bar: 20μm).





FIG 2 a and b - PHOTOMICROGRAPH SHOWING AT RIGHT 2 GRANULOMAS WITH CASEOUS NECROSIS IN THE CENTER, SURROUNDED BY EOSINOPHILIC AND FIBROBLASTS AND INFLAMMATORY CELLS (LYMPHOCYTES, NEUTROPHILS, HETEROPHILES AND MELANOMACROPHAGES) AND AN AREA OF CALCIFICATION (ARROW). HE. BAR = 200μm. ON THE LEFT, AT HIGHER MAGNIFICATIONS, A GRANULOMA AND NUMEROUS MELANOMACROPHAGES (BROWN). HE. (Bar: 50 μm).

It was observed, thus, lymphocytes, neutrophils and heterophiles. Some macrophages alone or in groups, were filled with golden-yellow substance (melanomacrophage) next to the granulomatous or degenerative lesions. The most severe changes were observed in the kidney that showed convoluted tubules in vacuolar degeneration or necrosis. Glomeruli were also visualized in degenerating, necrotic or deformed, hypo- or hyperplastic and presenting increased Bowman's space. Nephrocalcinosis was observed in 2 cases.

With the transmission electron microscope, *Mycobacterium* spp was also observed (Fig 3 a and b) in 8 fishes, these same animals that were positive for the Z-N technique.




FIG 3 a and b - ELECTRON MICROGRAPH OF MYCOBACTERIUM SPP IN FISH LIVER AND VISUALIZED BY NEGATIVE STAINING TECHNIQUE. AT THE ARROWS, THE TRIPLE LAYER ENVELOPE. (Bar: 140 nm)

Slow-growing scotochromogenic colonies were obtained from one fish (Fig 4). These colonies were catalase positive and did not reduce nitrate. The nearly entire 16S rDNA gene (1456pb) sequence obtained showed 99.65% identity with *M. gordonae* type strain (ATCC 14470). This was deposited in GenBank (accession number JN899581). We also sequenced the 16S rDNA gene of the type strain of *M. gordonae* ATCC 14470, (accession number JN899579) for comparative identification purposes, as in the sequence GenBank (X52923) had 8 unidentified (N) bases.




FIG 4 - COLONIES OF MYCOBACTERIUM GORDONAE ISOLATED FROM FISH IN PETRAGNANI MEDIUM

IV. DISCUSSION AND CONCLUSION

Brazil has enormous potential for animal and fish farming production, given its vast land, water sources and favorable weather conditions. In Brazil, fish are typically raised in lakes for consumption, although in this study they were not intended for consumption.

High concentrations of fish can favor the onset epizootic disease outbreaks caused by *Mycobacteria* spp, although in natural environmental conditions spontaneous disease outbreaks can also occur (GAUTHIER & RHODES, 2009; HECKERT et al., 2001; RAMSAY et al., 2009).

Mycobacteriosis can occur in several species of fish. In addition, some species of *Mycobacterium* spp. can be transmitted to humans by occupational or recreational source (BHATTY et al., 2000; GAUTHIER & RHODES, 2009; JACOBS et al., 2009; REAVILL & SCHMIDT, 2012).

In Brazil, there are a few studies on mycobacteriosis in ectotherms. MOK and CARVALHO in 1984, described the presence of *M. chelonae* in *Bufo marinus* and *B. granulosis*, and although mycobacteriosis outbreaks in frog farms have been reported by FERREIRA et al., 2006. In fish, was notification by ISHIKAWA et al., 2001 and ROMANO et al., 2012.

Histopathological examinations are important for early diagnosis of mycobacterial infection in fish. Granulomas are suggestive of mycobacteriosis but are not pathognomonic of the disease; acid-fast bacilli must be visualized in the lesions. A positive culture will provide a definitive diagnosis, but it is not very easy to isolate mycobacterias at 37° C, the optimum temperature for human pathogens. Fishes isolate are well-characterized by molecular methods.

In this study, we isolated acid-fast bacillus from one fish which were identified as *M. gordonae* by molecular methods. It is recommended that further, more in-depth, studies are undertaken to gain a better insight of the impact of this disease in cultured and wild fish species in Brazil.

ACKNOWLEDGEMENTS

The authors want to Thanks FAPESP

REFERENCES

- [1] A. Telenti, F. Marchesi, M. Balz, F. Bally, E.C. Bottger, T. Bodmer. Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis. *Journal of Clinical Microbiology* 31(2), 175–178, 1993.
- [2] C.E. Campos, P.C. Caldas, H. Ohnishi, T. Watanabe, K. Ohtsuka, S. Matsushima, N.V. Ferreira, M.V. Silva, P. Render, L.D. Carvalho, R.F. Medeiros, A.B.B.U.D. Filho, F.C. Montes, T.C. Galvão, J.P. Ramos. First isolation of *Mycobacterium kyorinense* from clinical specimens in Brazil. *Journal of Clinical Microbiology* 50 (7), 2477–2478, 2012.
- [3] C.M. Ishikwa, E.R. Matushima, C.W.O. Souza, J. Timenetsky, M.J. Ranzani-Paiva. Micobacteriose em peixes. *Boletim do Instituto de Pesca* 27, 231–242, 2001.
- [4] C.Q.F. Leite, C.W.O. Souza, S.R. Leite. Identification of Mycobacteria by Thin Layer Chromatographic Analysis of Mycolic Acids and Conventional Biochemical Method: Four Years of Experience. *Memórias do Instituto Oswaldo Cruz* 93(6), 801–805, 1988.
- [5] C.Y. Turenne, R. Wallace, M.A. Behr. *Mycobacterium avium* in the Postgenomic Era. *Clinical Microbiology Reviews* 20(2), 205–229, 2007.
- [6] D.R. Reavill, R.E. Schmidt. Mycobacterial lesions in fish, amphibians, reptiles, rodents, lagomorphs, and ferrets with reference to animal models. *The Veterinary Clinics of North America: Exotic Animal Practice* 15(1), 25–40, 2012.
- [7] D.T. Gauthier, M.W. Rhodes. Mycobacteriosis in fishes: A review. *The Veterinary Journal* 180(1), 33–47, 2009.
- [8] E. Bataillon, L. Dubard, L. Terre. Un nouveau type de tuberculose. *Comptes rendus des Séances de la Société de Biologie* 49, 446–449, 1987.
- [9] E. Tortoli. Impact of Genotypic Studies on Mycobacterial Taxonomy: the New Mycobacteria of the 1990s. *Clinical Microbiology Reviews* 16(2), 319–354, 2003.
- [10] F. Bier, M.L. Boschirolini, M.F. Thorel, L.A. Guilloteau. Zoonotic aspects of *Mycobacterium bovis* and *Mycobacterium avium*-intracellulare complex (MAC). *Veterinary research* 36(3), 411–436, 2005.
- [11] G.L. Fite, F.J. Cambre, M.H. Turner. Procedures for demonstrating lepra bacilli in paraffin sections. *Archives of Pathology*, V43, p624-625, 1947.
- [12] I.N. Kantor. Bacteriologia de la tuberculosis humana y animal. Serie de Monografías científicas Y Técnicas. Rev. Ied. Buenos Aires: Centro Panamericano de Zoonosis, 1998.
- [13] J.M. Grange, M.D. Yates. Infections caused by opportunist mycobacteria: a review. *Journal of the Royal Society of Medicine* 79(4), 226–229, 1986.
- [14] J.M. Jacobs, C.B. Stine, A.M. Baya, M.L. Kent. A review of mycobacteriosis in marine fish. *Journal of Fish Diseases* 32(2), 119–130, 2009.
- [15] J.M. Ramsay, V. Watral, C.B. Schreck, M.L. Kent. Husbandry stress exacerbates mycobacterial infections in adult zebrafish, *Danio rerio* (Hamilton). *Journal of Fish Diseases* 32(11), 931–941, 2009.
- [16] K.M. Astrofsky, M.D. Schrenzel, R.A. Bullis, R.M. Smolowitz, J.G. Fox. (2000). Diagnosis and management of atypical *Mycobacterium* spp. infections in established laboratory zebrafish (*Brachydanio rerio*) facilities. *Comparative Medicine* 50(6), 666 – 672, 2000.
- [17] L.A. Romano, L.A. Sampaio, M.B. Tesser. Micobacteriosis by *Mycobacterium marinum* in Brazilian flounder *Paralichthys orbignyanus* and barber goby *Elacatinus figaro*: histopathological and imunohistochemical diagnosis. *Pesquisa Veterinária Brasileira* 32(3), 254–258, 2012.
- [18] M.A. Bhatti, D.P.J. Turner, S.T. Chamberlain. *Mycobacterium marinum* hand infection: case reports and review of literature. *British Journal of Plastic Surgery* 53(2), 161–165, 2000.
- [19] M.A. Hayat, S.E. Miller. Negative Staining. Mc. Graw-Hill Publ. Company. 253 p. 1990.
- [20] P. Bemer-Melchior, H.B. Drugeon. Inactivation of *Mycobacterium tuberculosis* for DNA Typing Analysis. *Journal of Clinical Microbiology* 37(7), 2350–2351, 1999.
- [21] R. Ferreira, L.S. Fonseca, A.M. Afonso, M.G. Silva, M.H. Saad, W. Lilenbaun. A report of mycobacteriosis caused by *Mycobacterium marinum* in bullfrogs (*Rana catesbeiana*). *The Veterinary Journal* 171(1), 177–180, 2006.
- [22] R.A. Heckert, R.A. Eckert, S. Elankumaran, A. Milani, A. Baya. Detection of a New *Mycobacterium* Species in Wild Striped Bass in the Chesapeake Bay. *Journal of Clinical Microbiology* 39(2), 710–715, 2001.
- [23] S. Brenner, R.W. Horne. A negative staining method for high resolution electron microscopy of viruses. *Biochimica et Biophysica Acta* 34, 103, 1959.
- [24] T.M. Shinnick, R.C. Good. Mycobacterial taxonomy. *European Journal of Clinical Microbiology & Infectious Diseases* 13(11), 884–901, 1994.
- [25] W.Y. Mok, M. Carvalho. Occurrence and experimental infection of toads (*Bufo marinus* and *B. granulosis*) with *Mycobacterium chelonae* subs. *abscessus*. *Medical Microbiology* 18, 327-333, 1984.