

Evaluation of mass trapping for control of Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae) in Satsuma mandarin in Hatay province of Turkey

Nihat DEMİREL¹, Eda AKYOL²

Mustafa Kemal University, Faculty of Agriculture, Department of Plant Protection, 31034 Hatay-Turkey.

*Corresponding author: Dr. Nihat DEMİREL, Mustafa Kemal University, Faculty of Agriculture, Department of Plant Protection, 31034, Hatay-Turkey. [Tel: 90 (326) 245 58 45; Fax: 90 (326) 245 58 32; email: ndemirel@mku.edu.tr]

Abstract— The Mediterranean fruit fly (Medfly), *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae), is one of the most important pests of citrus in Turkey. The objective of this study was to evaluate mass trapping for the control of Medfly in Satsuma mandarin in Hatay province of Turkey. The studies were conducted in 2011-2012 using eostrap® invaginada traps baited with % 95 Trimedlure impregnated in a polymeric plug-type dispenser. In the first year, 48 traps per 0.7 ha were placed in an experimental site from 1st August to December. In the second year, 23 traps per 0.7 ha were placed in the same site from 14 August to December. After two years of the study, the population density of medfly varied in each of the sampling year. In the first year, a total of 8968 medfly adults were caught by traps. The largest mean of catches per trap were recorded on 31 October (64.21), followed by 24 October (31.29), 17 October (22.48), 7 November (20.64), 3 October (17.60) and 10 October (16.71). In the second year, a total of 1307 medfly adults were caught by traps. The largest mean of the catches per trap were recorded on 25 September (7.35), followed by 13 November (5.83), 6 November (5.52), 18 September (5.43) and 30 October (4.26), respectively. The percentages of damage rates of medfly observed in both years. The damage rates of Medfly were 10.91 and 8.56 % in 2011 and 2012, respectively. In conclusion, the population density of medfly on satsuma mandarin increased in September and October due to high temperature. The mass trapping was not enough to control medfly on satsuma mandarin. Therefore, the mass trapping should be used with pesticides to decrease the population density of medfly during September and October in Hatay province of Turkey.

Keywords— Medfly, trimedlure, traps, satsuma mandarin, Turkey.

I. INTRODUCTION

Satsuma mandarin, *Citrus unshiu* Marc., (Rutaceae: Sapindales) is one of the main cultivated *Citrus* variety in Turkey. Satsuma mandarin production is consisting of approximately 23,413.1 ha with a total produce of 795.050 tons of fruit per annum in Turkey, and Hatay province's share is 10,466.1 ha and 402.601 tons [1]. The Mediterranean fruit fly (Medfly), *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae) is the most destructive pest among economically important fruit fly species [2,3]. The medfly is a highly adaptive polyphagous tropical fruit fly attacking more than three hundred and fifty plant species [4,5]. The female flies lay eggs below the skin of fruits which are destroyed by larval feeding [6].

Protein bait sprays mixed with malathion or low toxicity insecticides, spinosad, lambda-cyhalothrin are successfully used to control medfly populations [7,8]. Traps baited with trimedlure are important tools for detection, monitoring and controlling of the medfly [2]. The mass trapping has proved to be effective in suppression of the Medfly and advantage of reduced environmental impact when comparing with toxic bait sprays [9,10]. In addition, mass trapping has been developed in several countries using traps baited with trimedlure, hydrolysed proteins or food-based attractants [11,12,13, 14, 15, 16,17]. This method has been conducted on different fruit trees like peach [18], cherimoya [19], citrus [14,20,21], fig [22], persimmon and coffee [7] and apple[23]. The purpose of the current study was to evaluate mass trapping for the control of Medfly in Satsuma mandarin in Hatay province of Turkey.

II. MATERIALS AND METHODS

The study was conducted in 2011-2102 at a satsuma mandarin orchard in Hatay province of Turkey. The study was carried out using the Eostrap® invaginada traps (Sanidad Agricola Econex, Santomera, Murcia, Spain) baited with % 95 Trimedlure, (formulated in a polymeric plug-type dispenser) (Sanidad Agricola Econex, Santomera, Murcia, Spain) and dichlorvos or 2.2- dichlorovinyl dimethyl phosphate (DDVP) tablet (Sanidad Agricola Econex, Santomera, Murcia, Spain).

In the first year, the study was conducted with 48 traps/0.7 ha and placed in the experimental site from 1st August to December. In the second year, the study was carried out with 23 traps/0.7 ha and placed in the same site from 14th August to December. The traps were placed 1.5 m above ground and checked weekly, trapped medfly adults were counted and removed from the traps. The trimedlure and DDVP tablet in traps were replaced with the new ones in every 90 days. The fruit damage assessment was measured by the percentage of medfly punctures during the harvest. For this purpose, all fruits from satsuma mandarin orchard were harvested and checked for medfly punctures and the percentage of the infested fruits were measured as weight at the packaging house.

III. RESULTS

The population density of *C. capitata* varied in each of the sampling year. In the first year, the mass trapping was conducted with 48 traps/0.7 ha in satsuma mandarin orchard and a total of 8968 medfly adults were caught by traps (Figure 1). The first adults were caught by the traps on 8 August, and the population density of this pest was gradually low from 8 August through 12 September, while it increased from 19 September to 31 October. The largest mean of catches per trap were recorded on 31 October (64.21), followed by 24 October (31.29), 17 October (22.48), 7 November (20.64), 3 October (17.60) and 10 October (16.71).




FIGURE 1. MEAN (±SE) CATCHES OF MEDFLY ADULTS IN TRAPS BAITED WITH TRIMEDLURE (1AUGUST–28 NOVEMBER, 2011) AT SATSUMA MANDARIN ORCHARD IN ANTAKYA DISTRICT




FIGURE 2. MEAN (±SE) CATCHES OF MEDFLY ADULTS IN TRAPS BAITED WITH TRIMEDLURE (15 AUGUST–11 DECEMBER, 2012) AT SATSUMA MANDARIN ORCHARD IN ANTAKYA DISTRICT

In the second year, the mass trapping was conducted with 23 traps/ha at the same orchard and a total of 1307 medfly adults were caught by traps (Figure 2). The population density of medfly was very low comparing with previous year. The first adults were caught by the traps on 21 August, and the population density of this pest was significantly high from 4 September to 25 September and 30 October through 13 November and yet it was significantly low on 2-23 October due to heavy rain in the sampling orchard. The largest mean of catches per traps were recorded on 25 September (7.35), followed by 13 November (5.83), 6 November (5.52), 18 September (5.43) and 30 October (4.26).

The percentage of the total caught over this period varied for each of the sampling month in both years. The distribution of the total caught of this pest according to the months as percentages were 1.17 (August), 6.55 (September), 71.32 (October) and 10.14 (November) in 2011 (Figure 3). These percentages were 3.67 (August), 32.13 (September), 16.37 (October) and 26.70 (December) in 2012 (Figure 4). The percentages of damage rates of medfly observed in both years. The damage ratios of medfly were recorded at 10, 91 percent in 2011, and 8, 56 percent in 2012 (Figure 5).




FIGURE 3. PERCENTAGE OF THE TOTAL MEDFLY ADULTS CAUGHT OVER THE SAMPLING PERIOD AT SATSUMA MANDARIN ORCHARD IN ANTAKYA DISTRICT.




FIGURE 4. PERCENTAGE OF THE TOTAL MEDFLY ADULTS CAUGHT OVER THE SAMPLING PERIOD AT SATSUMA MANDARIN ORCHARD IN ANTAKYA DISTRICT.




FIGURE 5. PERCENTAGE OF THE DAMAGED FRUITS BY MEDFLY IN SATSUMA MANDARIN ORCHARD IN ANTAKYA DISTRICT

IV. DISCUSSION

The mass trapping has shown significantly effective pest management tool for the Medfly and developed in several countries using traps baited with trimedlure, hydrolysed proteins or food-based attractants [2, 7, 9, 12, 13, 14, 15, 16, 17, 24, 25, 26, 27, 28, 29]. The synthetic food-based attractants, trimethylamine (TMA), ammonium acetate (AA) and putrescine (P) were the more appropriate for mass-trapping of the medfly [9, 11, 14, 15, 30,31, 32].

A various number of traps were used with mass trapping to control medfly on different host plants. Martinez-Ferrer et al. [10] suggested that 25 trap per ha (Maxitrap Model baited with Ferag. CC D TM® attractant) can be a good stand-alone control method against the Medfly in citrus groves in Spain. In addition, IAEA [2] suggested that 20-25 traps per ha density was required by using a mass trapping technique for medfly. Martinez-Ferrer et al. [10] reported that for the Clemenules variety, a 25 traps per ha were enough to capture adults flying within the grove and attract the foraging medflies, as low percentage of fruits were attacked. A number of researchers reported that if the population density of medfly increase, the number of trap per ha will be increased. Therefore, during September-October Medfly population is high, a 50 trap per ha density has been widely accepted as appropriate for citrus orchards [9,32,33,34,35,36,37]. Martinez-Ferrer et al. [10] reported that for the early-season varieties, 50 traps per ha captured as many adults as did 75 and 100 traps per ha, but not enough to diminish the adult medflies foraging in the grove under accepted levels.

Several studies were conducted by many researchers to evaluate the population density of medfly on various host plants. The present results indicated that the population density of medfly was high in September and October in both year. Our results are similar to those of [10,38,39] reported that the population density of medfly was significantly high in September and October due to high temperatures in citrus- producing area.

The percentages damage ratios of medfly varied on different host plants. Martinez-Ferrer *et al.* [37] found that on early-season varieties (Loretina and Marisol), mass trapping at a density of 50 Maxitrap® (Probodelt®) traps/ha baited with Ferag CC D TM® (SEDQ) and chemical treatments with Malafin® (malathion) or SpintorCebo® (spinosad) of the entire groves or the perimeter, can protect well the fruits from the Medfly attack as <2% were damaged in the harvest. However, on the mid-season variety (Clemenules), the adult population decline, and the number of traps could be lowered to only 25/ha and this tactic applied alone was efficient enough to obtain less than 0.5% of punctured fruits. Boulahia- Kheder *et al.* [40] reported that combination of 4 aerial spinosad sprays, sanitation and female mass trapping with Moskisan® + Biolure® Unipack, the damage on Navel oranges reached only 5% at harvest. However, Boulahia-Kheder *et al.*[41] reported that mass trapping as a single technique it wasn't enough to protect the Navel oranges with more than 30% of damaged fruits at the harvest.

V. CONCLUSION

The present study was conducted by traps baited with trimedlure to evaluate of mass trapping for the control of Medfly in Satsuma mandarin in Hatay province of Turkey. As a result of two-year investigations, the population density of this pest varied in each of the sampling year. In the first year, 48 trap per 0.7 ha were used with mass trapping to control of the medfly. A significant number of medfly were caught by traps and yet the medfly caused significant damages on satsuma mandarin due to high population density in September and October. In the second year, 23 trap per 0.7 ha were used with mass trapping to control of this pest. The population density of medfly was significantly low comparing with that of the previous year. However, the mass trapping was not enough to control this pest because of high population density

ACKNOWLEDGEMENTS

This project was supported by University of Mustafa Kemal of Scientific Research Projects (BAP) (project number: 1105 Y 0105 (191)).

REFERENCES

- [1] Anonymous, 2016. The summary of Agricultural Statistics (21.10.2017 version) <https://biruni.tuik.gov.tr/bitkiselapp/bitkisel.zul>.
- [2] International Atomic Energy Agency, 2003. Trapping guidelines for area-wide fruit fly programmes. International Atomic Energy Agency. Vienna, Austria. 47 p. Retrived from : http://www-pub.iaea.org/MTCD/Publications/PDF/TG-FFP_web.pdf
- [3] Thomas, M.C., Heppner, J.B., Woodruff, R.E., Weems, H.V., Steck, G.J., Fasulo, T.R. 2007. Mediterranean Fruit Fly. *Ceratitidis Capitata* (Wiedemann) (Insecta: Diptera: Tephritidae). University of Florida Ifas Extension. <http://edis.ifas.ufl.edu/pdffiles/IN/IN37100.pdf> Accessed on 13 October 2017.
- [4] Weems, H.V., Jr. 1981. Mediterranean fruit fly, *Ceratitidis capitata* (Wiedemann) (Diptera: Tephritidae). Entomology Circular, Division of Plant Industry, Florida Department of Agriculture and Consumer Services. 12 pp.
- [5] Liquido, N.J., Shinoda, L.A., and Cunningham, R.T. 1991. Host plants of the Mediterranean fruit (Diptera,Tephritidae) an annotated world review, Miscellaneous Publications 77. Entomol. Soc. Am., Lanham, MD. 1863-1878.
- [6] Christenson, L.D., and Foote, R.H. 1960. Biology of fruit flies. Annual Review of Entomology. 5: 171-192.
- [7] McQuate, G.T., Sylva, C.D., and Jang, E.B. 2005. Mediterranean fruit fly (Diptera: Tephritidae) suppression in persimmon through sprays in adjacent coffee plantings. Journal of Applied Entomology. 2: 110-117.

- [8] Urbaneja, A., Chueca, P., Monton, H., Pascual-Ruiz, S., Dembilio, O., Vanaclocha, P., Abad-Moyano, R., Pina, T., and Castanera, P. 2009. Chemical alternatives to malathion for controlling *Ceratitidis capitata* (Diptera: Tephritidae), and their side effects on natural enemies in Spanish citrus orchards. *J. Econ. Entomol.* 102: 144-151.
- [9] Navarro-Llopis, V., Alfaro, F., Domínguez, J., Sanchis, J. and Primo, J. 2008. Evaluation of traps and lures for mass trapping of Mediterranean fruit fly in citrus groves. *J. Econ. Entomol.* 101:126-131.
- [10] Martínez-Ferrer, M.T., Campos, J.M., and Fibla, J.M. 2010. Field efficacy of *Ceratitidis capitata* (Diptera: Tephritidae) mass trapping technique on clementine groves in Spain. *Journal of Applied Entomology.* 136: 181-190.
- [11] Epsky, N.D., Hendrichs, J., Katsoyannos, B.I., Vasquez, L.A., Ros, J.P., Zumreoglu, A., Pereira, R., Bakri, A., Seewooruthun, S.I. and Heath, R.R. 1999. Field evaluation of female-targeted trapping systems for *Ceratitidis capitata* (Diptera: Tephritidae) in seven countries. *J. Econ. Entomol.* 92: 156-164.
- [12] Katsoyannos, B.I., Papadopoulos, N.T., Heath, R.R., Hendrichs, J. and Kouloussis, N.A. 1999. Evaluation of synthetic food-based attractants for female Mediterranean fruit flies (Dipt., Tephritidae) in McPhail type traps. *J. Appl. Entomol.* 123: 607-612.
- [13] Cohen, H. and Yuval, B. 2000. Perimeter trapping strategy to reduce Mediterranean fruit fly (Diptera: Tephritidae) damage on different host species in Israel. *J. Econ. Entomol.* 93: 721-725.
- [14] Miranda, M.A., Alonso, R. and Alemany, A. 2001. Field evaluation of Medfly (Diptera, Tephritidae) female attractants in a Mediterranean agrosystem (Balearic Islands, Spain). *J. Appl. Entomol.* 125: 333-339.
- [15] Heath, R.R., Epsky, N.D., Midgarden, D. and Katsoyannos, B. 2004. Efficacy of 1,4-Diaminobutane (Putrescine) in a food based synthetic attractant for capture of mediterranean and mexican fruit fly (Diptera: Tephritidae). *J. Econ. Entomol.* 97:1126-1131.
- [16] Ortu, S., Lentini, A. and Cocco, A. 2005. Strategie di lotta per il contenimento di *Ceratitidis capitata* (Wied.) in agrumicoltura. *Informatore Fitopatologico.* 55: 28-34.
- [17] Alemany, A., Miranda, M.A., Alonso, R. and Escorza, C.M. 2006. Changes in the spatial and temporal population density of the Mediterranean fruit fly (Diptera: Tephritidae) in a citrus orchard. *Span. J. Agric. Res.* 4:161.
- [18] Sastre, C. 1999. Eficacia de los productos fitosanitarios en el control de la mosca de la fruta *Ceratitidis capitata* (Wied.). *Phytoma España.* 114: 75-77.
- [19] Ros, J. P., Escobar, I., García Tapia, F. J. and Aranda, G. 2000: Pilot experiment to control Medfly, *Ceratitidis capitata* (Wied.) (Diptera: Tephritidae) using mass trapping technique in a Cherimoyer (*Annona cherimola* Miller) orchard. In: Area-wide control of fruit fly and other insect pests (ed. K. H. Tan): 639-643. Penerbit University Sains Malaysia, Penang.
- [20] Alonso Muñoz, D., Soler, J. M., García Marí, F., and Blas Correas, M. 1999. Frutect®: control de la mosca de la fruta *Ceratitidis capitata* Wied. en el cultivo de los cítricos. *Levante Agrícola.* 347: 204-211.
- [21] Llorens, J. M., Vinaches, P., Marco, F. and Llorens, J. 2004: Control de *Ceratitidis capitata* Wied. using mass trapping with Tephri traps and Biolure-3 component Lure (Tripack) in fig trees. 5^o Meeting of the working group on fruit flies of the Western Hemisphere, 16-21 May, Florida. Abstract: 38.
- [22] Escudero, A., Vilajeliu, M. and Batllori, L. 2005. Captura masiva para el control de a mosca mediterránea de la fruta (*Ceratitidis capitata* Wied.) en manzano. *Phytoma España.* 171: 26-31.
- [23] Ortu, S. and Prota, R. 1988. Biotechnical control means adopted against *Ceratitidis capitata* Wied. in clementine groves. *IOBC/wprs Bull.* 11:14-19.
- [24] Avery, J.W., Chambers, D.L., Cunningham, R.T. and Leonhardt, B.A. 1994. Use of ceralure and trimedlure in Mediterranean fruit fly (Diptera: Tephritidae) mass-trapping tests. *J. Entomol. Sci.* 29:543-556.
- [25] Sastre C, Melo J.C. and Borrelli G. 1999. La captura de hembras: una posible salida en el control de la mosca de la fruta (*Ceratitidis capitata* Wied.) en melocotonero. *Phytoma.* 113: 42-46.
- [26] Ros, J.P., Gomila, J., Reurer, M., Pons, P. and Castillo, E. 2002. The use of mass trapping against Medfly (*Ceratitidis capitata* (Wied.)) in a sustainable agriculture system on Minorca Island, Spain. *Proceedings of the 6th International Symposium on fruit flies of economic importance, Stellenbosch, South Africa, 6-10 May 2002*, pp. 361-364.
- [27] Garcia, G., Wong, E., Marquez, A.L., Garcia, S., Olivero, J. and Garcia Mari, F. 2003. Evaluation and comparison of mass-trapping methods for the control of *Ceratitidis capitata* Wied., in citrus orchards. *Bull. OILB SROP.* 26:85.
- [28] Jemâa, J.M.B., Bachrouch, O., Allimi, E. and Dhouibi, M.H. 2010. Field evaluation of Mediterranean fruit fly mass trapping with Tripack® as alternative to malathion baitspraying in citrus orchards. *Spanish Journal of Agricultural Research.* 8: 400-408.
- [29] Ros, J.P., Castillo, E., Crespo, J., Latorre, Y., Martin, P., Miranda, M.A., Moner, P. and Sastre, C. 1997. Evaluación en campo de varios atrayentes sintéticos para la captura de hembras de la mosca mediterránea de la fruta *Ceratitidis capitata* Wied. (Díptera: Tephritidae). *Boletín de Sanidad Vegetal, Plagas.* 23: 393-402.
- [30] Heath, R.R., Epsky, N.D., Dueben, B.D., Rizzo, J. and Jeronimo, F. 1997. Adding methylsubstituted ammonia derivatives to a food-based synthetic attractant on capture of the Mediterranean and Mexican fruit flies (Diptera: Tephritidae). *J. Econ. Entomol.* 90: 1584-1589.
- [31] Alemany, A., Alonso, D. and Miranda, M.A. 2004. Evaluation of improved Mediterranean fruit fly attractants and retention systems in the Balearic Islands (Spain). Pages 355-359. In *Proceedings 6th International Symposium on Fruit Flies of Economic Importance, 6-10 May, 2004, Stellenbosch, South Africa.*
- [32] Primo, E. 2004. Plan Valenciano de actuación contra la mosca de la fruta (*Ceratitidis capitata* Wied.) campaña 2004. *Agricultura: Revista agropecuaria.* 867: 790-793.
- [33] Ros, J.P., Wong, E., Olivero, J., Rubio, J.R., Ma´rquez, A.L., Castillo, E. and Blas, P. 2005. Desarrollo de atrayentes y mosqueros para su integración en los programas de trampeo masivo contra la mosca de la fruta (“*Ceratitidis capitata*” Wied.) y la del olivo (“*Bactrocera oleae*” Gmel). *Bol. Sanid. Veg. Plagas.* 31: 599-607.

-
- [34] Navarro-Llopis, V., Sanchis-Cabanes, J., Ayala, I., Castana Giner, V. and Primo-Yufera, E. 2004. Efficacy of lufenuron as chemosterilant against *Ceratitidis capitata* in field trials. *Pest Manag. Sci.* 60: 914–920.
- [35] Leza, M.M., Juan, A., Capllonch, M. and Alemany, A. 2008. Female-biased mass trapping vs. bait application techniques against the Mediterranean fruit fly, *Ceratitidis capitata* (Dipt.: Tephritidae). *Journal of Applied Entomology* 132: 753–761.
- [36] Martinez-Ferrer, M.T., Campos, J.M. and Fibla, J.M. 2012. Field efficacy of *Ceratitidis capitata* (Diptera: Tephritidae) mass trapping technique on clementine groves in Spain. *Journal of Applied Entomology.* 136:181-190.
- [37] Martinez-Ferrer, M.T., Campos, J.M. and Fibla, J.M. 2006. Population dynamics of *Ceratitidis capitata* on citrus in northeastern Spain: influence of adjacent host fruit trees. *IOBC/WPRS Bull.* 29: 77–84.
- [38] Martinez-Ferrer, M.T., Alonso Muñoz, A., Campos Rivela, J.M., Fibla Queralt, J.M. and Garcia-Mar, F. 2007. Dinámica poblacional de la mosca de la fruta *Ceratitidis capitata* en tres zonas cítricas mediterráneas. *Levante Agrícola, Num.* 385: 92–98.
- [39] Boulahia-Kheder, S., Trabelsi, I. and Aouadi, N. 2012. From chemicals to IPM against the Mediterranean fruit fly *Ceratitidis capitata*. In Larramendy, M.L. and Soloneski, S. (eds.) *Integrated Pest Manage. Pest Control, Curr. Future Tactics.* Croatia: In Tech.
- [40] Boulahia-Kheder, S., Jerraya, A., Fezzani, M. and Jrad, F. 2010. First results in Tunisia on the mass-trapping an alternative way to control the Mediterranean fruit fly *Ceratitidis capitata* (Diptera: Tephritidae). *Annals INRAT.* 82: 168-180.