

Effect of interaction between different plant growth regulators on in vitro shoot multiplication of *Citrus latifolia* Tan. (persian lime)

Firoozeh Chamandoosti

Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO),
Tehran, Iran

PhD of Cellular and Developmental Biology, Assistant Professor of Iranian Research Institute of Plant Protection
Department of Plant Diseases

Abstract— In this paper a shoot multiplication is described for *Citrus latifolia* Tan. (persian lime) using nodal segment explants of young one – old – year trees by two different pathways contain with and without callusing phase. The best result for multiple shoot formation and regenerated shoot formation was 3.2 and 2.6 shoots per explants with 4.44 μM BA plus 0.053 μM NAA and 4.44 μM BA plus 0.049 μM IBA respectively. Alike shoot regeneration, shoot elongation was occurred in medium with 4.44 μM BA and 0.049 μM IBA. Micropropagated and regenerated plants are under other experiments.

Abbreviation: BA – 6 benzylaminopurine; IBA – Indole acetic acid; NAA – Naphtalene acetic acid; PGRs – Plant Growth Regulators.

Key Words: Persian lime; plant growth regulators; shoot multiplication.

I. INTRODUCTION

Accordant worldwide *Citrus* species are the most widely grown fruit crops in Iran. They contain vitamin C that is very useful for human nutrition. Also their fruits are important source of volatile oils, limonene, α -terpinene, β -terpinene, citral cumarins, bioflavonoids, vitamins, and mucilage (Rathore *et al.*, 2006). Beside apples and bananas, *Citrus* fruits are the most important fruit crops (FAO 2001). Also it is clear that the sustainable development of the *Citrus* industry is mainly dependent on a continuous supply of new and improved cultivars (Perez – Tornero *et al.*, 2010). For the *Citrus* industry to improve fruit quality and reduce biotic and abiotic stresses are major breeding objectives at any time (Wenwu *et al.*, 2007). *Citrus* varieties are propagated by both sexual and asexual methods. Generally, rootstocks are propagated sexually through seeds, while most of the commercial varieties are propagated by various asexual methods (Chaudhary 1994). These conventional techniques are also not free from risk of perpetuating in-born pathogens. However in vitro micropropagation technology can overcome some constraints to *Citrus* improvement and cultivation, and can increase fruit quality and resistance to diseases and environmental stresses (Gresser 1994). Also micropropagation systems with high multiplication rates are not only an important asexual method that can be used for the production of clonal plants, but also form the basis for the introduction of genetic variation by genetic transformation or mutagenesis. The genetic transformation of *Citrus* has been widely studied as a tool to generate transgenic plants with enhanced tolerance of biotic (Cardoso *et al.*, 2010; He *et al.*, 2011 and Ali *et al.*, 2012) and/or abiotic stresses (Bunnag and Tangpong 2012). In both cases, is necessary to be able to regenerate viable shoots, which can be propagated, by either organogenesis or somatic embryogenesis (Perez – Tornero *et al.*, 2010).

In this paper a protocol for cloning of *Citrus latifolia* Tan. (persian lime) using nodal explant was described. It should be noted that according to Iranian researchers that used graft inoculation and PCR methods, persian lime was tolerant to Candidatus *Phytoplasma aurantifolia*. Candidatus *Phytoplasma aurantifolia* is a serious threat to lime and other susceptible *Citrus* trees in southern Iran (Salehi *et al.*, 2005).

II. MATERIAL AND METHODS

This study was conducted in Iranian Research Institute of Plant Protection Department of Plant Diseases, Tehran – Iran

Young trees of *Citrus latifolia* Tan. (persian lime) were collected from Jahrom *Citrus* nursery, Jahrom – Fars – Iran

2.1 Surface sterilization and plant material preparation

One – old – year young tree persian lime (*Citrus latifolia* Tan.) were used as the source of explants from their nodal segments. 30 – 35 – old – day new shoots measuring 12 – 15 cm in length with 4 – 6 nodes were cut and collected in plastic bag and transferred to the laboratory from greenhouse for experiments.

2.2 Preparation and sterilization of explants

After defoliation of shoots, and cutting 0.5 – 0.7 cm in length nodal segments (explants) surface sterilized under laminar air flow with dipping in 25% hypochlorite sodium for 5 min. Then were rinsed in distilled water 4 – 5 times.

2.3 Culture media

Culture media consisted of MS (Murashig and Skoog 1962) salts and vitamins plus 3% sucrose that were solidified with 0.75 % agar agar. Also the media were supplemented with BA (0 – 8.88 μM), 0.053 μM NAA and 0.049 μM IBA. The pH of media was adjusted to 5.8 before gelling with 1N NaOH or HCl and after gelling autoclaved for 20 min at 121°C. Then the media dispensed into 9 – cm diameter petri dishes. The culture was incubated at $25 \pm 2^\circ\text{C}$ and under 16 – h photoperiod.

2.4 Experimental design and data analysis

Experiments were conducted in a completely randomized design with 4 replicates and 5 explants per replicate. The mean number of multiplied shoots per explants and the mean length of multiplied shoots assay carried out on MS media with 12 combinations of cytokinin (BA) and auxins (NAA and IBA). Data were analysed by Duncan Multiple Range Test.

III. RESULTS AND DISCUSSION

Shoot multiplication in the presence of media with different concentrations of BA and 0.053 μM NAA and or 0.049 μM IBA compared after nearly 4 weeks of culture. It can be seen that the highest number of multiplied shoots (3.2) per explants was observed in the media containing 4.44 μM BA and 0.053 μM NAA (Fig. 1 and Fig. 5). Also the highest length of shoot (10 cm) was obtained on medium with 4.44 μM BA and 0.049 μM IBA (Fig. 3 and Fig. 6). It is interesting that in these study observations such as shoot multiplication and growth of shoot (longitudinal growth and foliation) were affected by interaction between different plant growth regulators more rather than plant growth regulators solely. There are reports about positive effect of increasing concentration of BA on shoot multiplication. For example Mehdi Farshad and coworkers in 2014 resulted that BAP alone (from 8.8 μM to 26.6 μM) was significantly effective on shoot multiplication in *Chlorophytum borivilianum*. However there are reports that show shoot proliferation decreased with increasing concentration of BA alone (Komal *et al.*, 2013). This experience is in accordance with ours that frequency of response from nodal explants decreased with a progressive increase in the level of BA [8.88 and 4.44 μM BA, 4.75 – 8.20 (cm for length of shoots)] respectively.

As was mentioned earlier, the interaction between plant growths regulators have very important effects on induction of shoots also number of shoot multiplied. So as was shown in Fig. 2 the mean number of multiplied shoots on explants increased when BA was used solely (1.2 – 2.66 for 0.044 – 8.88 μM BA). The positive effect of BA on the induction of regenerated shoot is clear. For example the positive effect of BA on shooting in *Cucurbita maxima* Duch (Lee *et al.*, 2003), *Ruta graveolunse* L. (Ahmad *et al.*, 2010), tomato (*Lycopersicon esculentum* L.) (Rai *et al.*, 2012) and Japanese pear (Kadota *et al.* 2001). These results were somewhat similar to our results about the mean number of multiplied shoots. However it seams 4.44 μM BA either in terms of nature or in terms of level is suitable for shoot multiplication in *Citrus* genus. So that Al-Khayri and Al-Bahrany in 2001 expressed that Best results for multiple shoot formation, 8 shoots per node, were obtained with 4.44 μM BA and 2.32 μM kinetin.

Another interesting result of this study is the circumstance of shoot multiplication. As is clear in figures multiplication shoot in the presence of BA and NAA was direct while multiplication shoot in the presence of BA and IBA was indirect and accompanied by callusing phase (Fig.7 and Fig. 8). So the role of interaction between plant growth regulators in in vitro culture of plants is proved again.

Very different and important roles are demonstrated for plant cell, tissue and organ culture. Briefly shoot multiplication and shoot regeneration are two different kind of organogenesis with two different utilizations. This result implies that rapid plant regeneration system which could be used for the somaclonal variation induction are possible for persian lime, on the other hand in vitro propagation through lateral buds proliferation that is an efficient method for large scale production of true – to – type planting material of important plant (Doo and Iyyakkannu 2016) is practicable for this valuable plant. It is worth noting that for assessment the level of background genetic changes resulting from the tissue culture processes (Munthali *et al.* 1996) and to verify the “true-to-type” genotype of micropropagated plants with shoot multiplication and or shoot regeneration DNA-based marker techniques such as RFLPs (Nelke *et al.*, 1993) and or RAPD (Munthali *et al.*, 1996) are necessary (Komal *et al.*, 2013)




FIG 1. The mean number of multiplied shoot on medium with 8.88 – 0.044 μM BA and 8.88-0.044 μM BA plus 0.053 μM NAA




FIG 2. The mean number of multiplied shoot on medium with 8.88 – 0.044 μM BA and 0.049 μM IBA




FIG 3. The mean length of multiplied shoot on medium with 8.88 – 0.044 μM BA and 8.88 – 0.044 μM BA and 0.049 μM NAA




FIG 4. The mean length of multiplied shoot on medium with 8.88 – 0.044 μM BA and 0.049 μM IBA

Mean followed by same letter(s) are not significantly different



FIGURES 5 – 8. 5 – Multiplied shoot on medium with 4.44 μM BA and 0.053 μM NAA. 6 – Elongation of multiplied shoot on medium with 4.44 μM BA and 0.049 μM IBA. 7 and 8 - Multiplied shoot on medium with 0.44 μM BA and 0.049 μM IBA

REFERENCES

- [1] Ahmad N. Faisal M. Anis M. and Aref I.M. 2010. In Vitro callus induction and plant regeneration from leaf explants of *Ruta graveolense* L. South Africa Journal. 76(3): 597 – 600
- [2] Al-Khayri J.M. and Al-Bahrany A.M. 2001. In vitro micropropagation of *Citrus aurantifolia* (lime). Current Science. 81(9): 1242 – 1246
- [3] Ali S. Mannan A. Oirdi M.E. Waheed A. and Mirza B. 2012. *Agrobacterium*-Mediated Transformation of Rough Lemon (*Citrus jambhiri* Lush) with Yeast HAL2 Gene. BMC Research Notes. 5: 285
- [4] Bunnag S. and Tangpong, D. 2012. Genetic Transformation of *Citrus sinensis* L. with an Antisense ACC Oxidase Gene. American Journal of Plant Sciences. 3: 1336 – 1340
- [5] Cardoso S.C. Barbosa-Mendes J.M. Boscarol-Camargo R.L. Christiano R.S.C. Filho A.B. Vieira M.L.C. Mendes B.M.J. and Mourao Filho F.A.A. 2010. Transgenic Sweet Orange (*Citrus sinensis* L. Osbeck) Expressing the Attacin A Gene for Resistance to *Xanthomonas citri* subsp. *citri*. Plant Molecular Biology Reports. 28: 185 – 192
- [6] Chaudhary M. I. 1994. Fruit Crops. In: Malik M.N. (ed.), Horticulture, 422. National Book Foundation, Islamabad
- [7] Doo H. K. Iyyakkannu S. 2016. Influence of benzyladenine and thidiazuron on shoot regeneration from leaf and shoot tip explants of *Sedum sarmentosum* Bunge. Biological and Applied Sciences. 59: 1 – 6
- [8] FAO 2001. <http://apps.fao.org/lim500/nph-wrap.pl>.
- [9] Gresser J.W. 1994. In vitro culture of tropical fruits, in plant Cell and tissue culture. Edited by I K Vasil and T A Thorpe (Kluwer Academic Publishers, Dordrecht, The Netherlands). 475 – 496
- [10] He Y. Chen S. Peng A. Zou X. Xu L. Lei T. Liu X. and Yao L. 2011. Production and Evaluation of Transgenic Sweet Orange (*Citrus sinensis* Osbeck) Containing Bivalent Antibacterial Peptide Genes (Shiva A and Cecropin B) via a Novel *Agrobacterium*-Mediated Transformation of Mature Axillary Buds. Scientia Horticulturae. 18: 99 – 107
- [11] Kadota M. Lmizu K. and Hirano T. 2001. Double-phase in vitro culture using sorbitol increases shoot proliferation and reduces hyperhydricity in Japanese pear. Scientia Horticulturae. 89(3): 207 – 215
- [12] Komal G. Sharma R. Singh P. K. and Govind S. 2013. Micropropagation of seedless lemon (*Citrus limon* L. cv. Kaghzi Kalan) and assessment of genetic fidelity of micropropagated plants using RAPD markers. Physiology and Molecular Biology of Plants. 19(1): 137 – 145
- [13] Lee Y.K. Chung W. Hiroshi E. 2003. Efficient plant regeneration via organogenesis in winter squash (*Cucurbita maxima* Duch.). Plant Science 164: 413 – 418
- [14] Mehdi Farshad A. Maheran A.A Nurashikin K. and Ismanizan I. 2014. Effect of cytokinin types, concentrations and their interactions on in vitro shoot regeneration of *Chlorophytum borivillanum* Sant. & Fernandez. Electronic Journal of Biotechnology. 17: 275 – 279
- [15] Munthali M.T. Newbury H.J. Ford-Llyod B.V. 1996. The detection of somaclonal variants of beets using RAPD. Plant Cell Reports. 16: 474 – 478
- [16] Murashige T. and Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Phiol. Plant. 15: 473 – 497
- [17] Nelke M. Nowak I. Wright J.M. McLean N.L. 1993. DNA-based marker techniques to detect somaclonal variations. Plant Cell Reports. 13: 72–78
- [18] Perez-Tornero O. Tallon C. I. Porras I. 2010. An efficient protocol for micropropagation of lemon (*Citrus limon*) from mature nodal segments. Plant Cell Tissue and organ Culture. 100: 263 – 271
- [19] Rai G.K. Rai N.P. Kumar S. yadav A. and Rathaur S. 2012. Effects of explant age, germination medium, pre-culture parameters, inoculation medium, pH, washing medium, and selection regime on *Agrobacterium*-mediated transformation of tomato. In Vitro Cellular and Developmental Biology. 48: 565 – 578
- [20] Rathore J.S. Rathore M.S. Singh M. Singh R.P. and Shekhawat N. S. 2006. Micropropagation of mature tree of *Citrus limon*. Indian Journal of Biotechnology. 6: 239 – 244
- [21] Salehi M. Nejat N. Tavakoli A.R. and Izadpanah K. 2005. Reaction of Citrus cultivars to Candidatus *Phytoplasma aurantifolia* In Iran. Iranian Journal of Plant Pathology. 41(3): 363 – 376
- [22] Wenuu G. Dingli L. and Yanxin D. 2007. Citrus Transgenic: Current Status and Prospects. Transgenic Plant Journal. 1(1): 202 – 209.