

Exploring Orchid Nutrition: Fungal Associations, Atmospheric Absorption, and Root Western Ghats in Wayanad, India

Mr. Sabu V.U

Vayalarikil (H), Kalathuvayal (PO), Ambalavayal. 673593, Wayanad

*Corresponding Author

Received:- 08 January 2025/ Revised:- 14 January 2025/ Accepted:- 21 January 2025/ Published: 31-01-2025

Copyright @ 2025 International Journal of Environmental and Agriculture Research

This is an Open-Access article distributed under the terms of the Creative Commons Attribution

Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted

Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— Orchids (*Orchidaceae*) are a diverse plant family known for their unique nutritional adaptations, allowing them to thrive in a variety of ecosystems. This five-year study explores the intricate nutrient acquisition strategies of orchids in the Western Ghats, particularly in the Wayanad region, a biodiversity hotspot with a rich diversity of epiphytic and terrestrial orchid species. The research investigates five primary nutritional pathways: mycorrhizal symbiosis, atmospheric nutrient absorption, rainwater uptake, organic matter decomposition, and specialized root structures. A critical aspect of orchid survival is their reliance on mycorrhizal fungi, particularly during early developmental stages, where fungal partners provide essential nutrients in the absence of endosperm. Additionally, epiphytic orchids demonstrate the ability to absorb nitrogen and carbon dioxide from the atmosphere, supplementing their nutrient intake. Rainwater serves as a primary source of hydration and dissolved minerals, particularly in the high-humidity environments of the Western Ghats. Orchids growing in terrestrial conditions benefit from organic matter decomposition, utilizing nutrients released from decaying leaves and animal residues. Specialized root structures, such as velamen-covered aerial roots, enhance water and nutrient absorption efficiency.

This study, conducted by an experienced plant conservation fellow, combines field observations, laboratory analyses, and greenhouse experiments to provide practical conservation insights. Findings highlight the importance of preserving natural orchid habitats, maintaining fungal diversity, and understanding climate-related challenges to orchid survival. These insights contribute to orchid conservation efforts, sustainable propagation techniques, and ecological research on plant-fungi interactions in tropical environments.

Keywords— Orchid nutrition, mycorrhizal fungi, epiphytic adaptation, rainwater uptake, airborne nutrients, root adaptations, organic matter decomposition, Western Ghats, Wayanad biodiversity, plant conservation.

I. INTRODUCTION

Orchids (*Orchidaceae*) represent one of the largest and most diverse plant families, with over 25,000 species distributed across various ecosystems worldwide. They are particularly well-adapted to different climatic conditions, ranging from tropical rainforests to temperate and alpine regions. The Western Ghats, a UNESCO World Heritage site, is home to a significant number of orchid species, many of which are endemic and play crucial ecological roles in their habitats. The unique environmental conditions of this region, including high humidity, varied elevation, and dense forest cover, contribute to the remarkable diversity of orchids.

Within the Western Ghats, the Wayanad region stands out as an important hotspot for orchid diversity. The area is characterized by tropical evergreen and semi-evergreen forests, grasslands, and montane ecosystems that provide ideal conditions for both epiphytic and terrestrial orchids. Epiphytic orchids, which grow on trees without drawing nutrients from their hosts, rely heavily on atmospheric moisture, rainwater, and nutrient-rich organic debris. Terrestrial orchids, on the other hand, thrive in the forest understory, where they depend on soil nutrients and symbiotic mycorrhizal fungi for growth and survival.

Despite their ecological importance, orchids in the Wayanad region face numerous threats, including habitat destruction, climate change, and overcollection for commercial purposes. Understanding their nutritional strategies is crucial for developing effective conservation measures. This study, conducted over five years, aims to investigate the various environmental factors influencing orchid nutrition, with a particular focus on their reliance on mycorrhizal fungi, atmospheric nutrient absorption,

rainwater uptake, and organic matter decomposition. By examining these mechanisms, this research provides valuable insights into orchid ecology and proposes practical conservation strategies to ensure the survival of these vulnerable species in their native habitat.

The findings of this study have significant implications for plant conservation efforts, particularly in the context of biodiversity preservation in the Western Ghats. The results will contribute to a deeper understanding of orchid survival strategies and inform future policies and conservation initiatives aimed at protecting these delicate and ecologically significant plants.

II. METHODOLOGY

This longitudinal study was conducted across various elevations in Wayanad, including tropical evergreen forests and montane regions. The study involved field observations, controlled greenhouse experiments, and laboratory analyses of root structures, fungal associations, and nutrient uptake patterns. Key methods included:

- **Mycorrhizal analysis:**
- **Airborne nutrient quantification:**
- **Rainwater nutrient composition.**
- **Organic matter decomposition studies:**
- **Root absorption efficiency**

The study methodologies with examples of common wild orchids found in the Wayanad region of the Western Ghats.

TABLE 1
EXAMPLES OF COMMON WILD ORCHIDS FOUND IN THE WAYANAD REGION

Methodology	Description	Common Wild Orchids in Wayanad	Findings & Significance
Mycorrhizal Analysis	DNA sequencing and microscopy to identify fungal partners aiding orchid nutrition.	<i>Vanda tessellata</i> , <i>Dendrobium macraei</i>	Strong symbiotic relationships with mycorrhizal fungi, crucial for seed germination.
Airborne Nutrient Quantification	Measuring nitrogen and carbon dioxide absorption rates at different altitudes.	<i>Oberonia ensiformis</i> , <i>Aerides ringens</i>	Epiphytic orchids at higher altitudes showed enhanced airborne nutrient absorption.
Rainwater Nutrient Composition	Analyzing minerals and trace elements in rainwater absorbed by orchids.	<i>Pholidota imbricata</i> , <i>Rhynchostylis retusa</i>	Orchids efficiently utilize rainwater as a primary nutrient source in high humidity regions.
Organic Matter Decomposition Studies	Assessing the role of decomposing leaves, animal droppings, and dead plant material in nutrition.	<i>Habenaria longicorniculata</i> , <i>Goodyera procera</i>	Terrestrial orchids benefit from decayed organic matter and microbial interactions.
Root Absorption Efficiency	Evaluating specialized root structures under varying soil compositions and humidity conditions.	<i>Bulbophyllum neilgherrense</i> , <i>Anoectochilus roxburghii</i>	Presence of velamen layers enhances water and nutrient retention, aiding survival in different microhabitats.

III. RESULTS AND DISCUSSION

3.1 Mycorrhizal Fungi and Orchid Nutrition:

Our findings confirm that orchids in the Wayanad region depend significantly on mycorrhizal fungi, particularly during seed germination. Field studies revealed that certain fungal species were more prevalent at different elevations, indicating a strong ecological connection between orchid survival and fungal diversity in the Western Ghats. This symbiotic relationship ensures that orchids can access nutrients in nutrient-poor environments, a trait shared with many common wild orchids found in other tropical and temperate regions. However, differences exist in the specificity of fungal associations, with orchids in Wayanad showing a higher dependency on endemic fungal species compared to generalist fungi that associate with common wild orchids.

3.2 Airborne Nutrients as a Supplementary Source:

Orchids demonstrate the ability to absorb nitrogen and carbon dioxide from the air through their leaves. The study observed that epiphytic orchids, often found in the upper canopy of Wayanad's forests, were highly efficient in utilizing atmospheric nutrients due to their exposure to mist and humidity. In comparison, common wild orchids, especially those in temperate zones, exhibit lower efficiency in absorbing airborne nutrients due to drier atmospheric conditions and reduced humidity levels.

3.3 Role of Rainwater in Orchid Hydration and Nutrition:

Rainwater was found to be a primary water source, carrying dissolved minerals essential for orchid nutrition. Wayanad's high rainfall provided ample hydration, but seasonal variations influenced nutrient availability, affecting orchid growth cycles. Common wild orchids, particularly those found in arid and temperate climates, rely more on dew and occasional rainfall for hydration, often exhibiting drought-resistant adaptations.

3.4 Organic Matter and Nutrient Cycling:

Accumulated organic debris such as fallen leaves and animal droppings significantly contributed to the nutrient supply of terrestrial orchids. Decomposing plant matter enriched the forest floor, allowing orchids to extract essential nutrients through their roots and symbiotic fungi. Compared to common wild orchids, which also utilize organic matter for nutrition, Wayanad's orchids benefit from a more consistent supply of decomposing organic material due to dense forest coverage and high humidity levels.

3.5 Specialized Root Adaptations for Nutrient Uptake:

Orchid roots in the Wayanad region exhibited specialized adaptations, including velamen, a multi-layered epidermis that enhances water and nutrient absorption. Orchids in drier areas developed thicker velamen layers, while those in consistently humid regions had more permeable structures for rapid water uptake. Common wild orchids in drier regions have similarly thickened velamen layers, whereas those in temperate climates may develop finer root structures with increased branching to maximize nutrient uptake.

TABLE 2

COMPARISON OF NUTRITIONAL STRATEGIES BETWEEN WAYANAD ORCHIDS AND COMMON WILD ORCHIDS

Nutritional Strategy	Orchids in Wayanad	Common Wild Orchids
Mycorrhizal Fungi Dependency	High dependency, particularly on endemic fungal species	Moderate dependency, often with more generalist fungal associations
Airborne Nutrient Absorption	Highly efficient, especially among epiphytic orchids	Less efficient due to drier conditions in non-tropical regions
Rainwater Utilization	Primary water source, seasonal variations influence growth	Relies on dew and occasional rain, more drought-resistant adaptations
Organic Matter Utilization	High due to dense forest cover and continuous organic debris decomposition	Moderate, depending on habitat type (forests vs. open fields)
Root Adaptations	Velamen structure varies by humidity levels; thicker in drier areas	Thick velamen in arid conditions; finer, branched roots in temperate zones

This comparative analysis highlights the unique adaptations of orchids in the Wayanad region and their responses to environmental factors, setting them apart from common wild orchids found in diverse ecosystems worldwide. These findings provide valuable insights into conservation strategies, emphasizing the need to protect mycorrhizal fungi and maintain ecological conditions that support orchid survival.

IV. CONCLUSION

The results of this five-year study underscore the complex nutritional strategies orchids employ for survival in the Wayanad region of the Western Ghats. The reliance on mycorrhizal fungi, atmospheric nutrient absorption, rainwater uptake, and organic matter decomposition highlights their adaptability to various ecological conditions. These findings emphasize the need for

conservation efforts that prioritize habitat preservation and fungal diversity protection, ensuring the long-term survival of orchids in their native environments.

Moreover, this study highlights the importance of an integrated conservation approach that combines habitat protection, sustainable collection practices, and the promotion of public awareness regarding orchid conservation. The intricate relationship between orchids and their surrounding ecosystem reinforces the necessity of preserving both plant and microbial biodiversity.

Future research should explore the impact of climate change on orchid nutrient acquisition and assess potential applications of these findings in sustainable horticulture. By implementing the conservation strategies proposed in this study, we can help safeguard these ecologically significant plants and contribute to the broader goal of biodiversity conservation in the Western Ghats.

V. FUTURE RESEARCH DIRECTIONS

Further studies are recommended to explore the molecular mechanisms governing orchid-mycorrhizal interactions, the impact of climate change on nutrient uptake, and potential applications of these findings in horticulture and conservation biology.

REFERENCES

- [1] Arditti, J. (1992). *Fundamentals of Orchid Biology*. Wiley.
- [2] Benzing, D. H. (2000). *Bromeliaceae: Profile of an Adaptive Radiation*. Cambridge University Press.
- [3] Cameron, K. M., Chase, M. W., & Whitten, W. M. (1999). A phylogenetic analysis of the Orchidaceae. *American Journal of Botany*, 86(2), 208-224.
- [4] Dressler, R. L. (1993). *Phylogeny and Classification of the Orchid Family*. Cambridge University Press.
- [5] Hadley, G. (1982). *Orchid Mycorrhiza*. CRC Press.
- [6] Johnson, S. D., & Edwards, T. J. (2000). The structure and function of orchid pollinaria. *Annals of Botany*, 85(2), 267-273.
- [7] Rasmussen, H. N. (2002). Recent developments in the study of orchid mycorrhiza. *Plant and Soil*, 244(1-2), 149-163.
- [8] Smith, S. E., & Read, D. J. (2008). *Mycorrhizal Symbiosis*. Academic Press.
- [9] Swarts, N. D., & Dixon, K. W. (2009). Terrestrial orchid conservation in the age of extinction. *Annals of Botany*, 104(3), 543-556.
- [10] Zotz, G., & Winkler, U. (2013). Aerial roots of epiphytic orchids: the velamen radicum and its role in water and nutrient uptake. *Oecologia*, 171(2), 733-741.