



International Journal of

Environmental & Agriculture Research

www.ijoeear.com

ISSN
2454-1850



Volume-10, Issue-1, January 2024

Preface

We would like to present, with great pleasure, the inaugural volume-10, Issue-1, January 2024, of a scholarly journal, *International Journal of Environmental & Agriculture Research*. This journal is part of the AD Publications series in the field of *Environmental & Agriculture Research Development*, and is devoted to the gamut of Environmental & Agriculture issues, from theoretical aspects to application-dependent studies and the validation of emerging technologies.

This journal was envisioned and founded to represent the growing needs of Environmental & Agriculture as an emerging and increasingly vital field, now widely recognized as an integral part of scientific and technical investigations. Its mission is to become a voice of the Environmental & Agriculture community, addressing researchers and practitioners in below areas.

Environmental Research:

Environmental science and regulation, Ecotoxicology, Environmental health issues, Atmosphere and climate, Terrestrial ecosystems, Aquatic ecosystems, Energy and environment, Marine research, Biodiversity, Pharmaceuticals in the environment, Genetically modified organisms, Biotechnology, Risk assessment, Environment society, Agricultural engineering, Animal science, Agronomy, including plant science, theoretical production ecology, horticulture, plant, breeding, plant fertilization, soil science and all field related to Environmental Research.


Agriculture Research:

Agriculture, Biological engineering, including genetic engineering, microbiology, Environmental impacts of agriculture, forestry, Food science, Husbandry, Irrigation and water management, Land use, Waste management and all fields related to Agriculture.

Each article in this issue provides an example of a concrete industrial application or a case study of the presented methodology to amplify the impact of the contribution. We are very thankful to everybody within that community who supported the idea of creating a new Research with *IJOEAR*. We are certain that this issue will be followed by many others, reporting new developments in the Environment and Agriculture Research Science field. This issue would not have been possible without the great support of the Reviewer, Editorial Board members and also with our Advisory Board Members, and we would like to express our sincere thanks to all of them. We would also like to express our gratitude to the editorial staff of AD Publications, who supported us at every stage of the project. It is our hope that this fine collection of articles will be a valuable resource for *IJOEAR* readers and will stimulate further research into the vibrant area of Environmental & Agriculture Research.



Mukesh Arora
(Managing Editor)



Dr. Bhagawan Bharali
(Chief Editor)

Fields of Interests

Agricultural Sciences	
Soil Science	Plant Science
Animal Science	Agricultural Economics
Agricultural Chemistry	Basic biology concepts
Sustainable Natural Resource Utilisation	Management of the Environment
Agricultural Management Practices	Agricultural Technology
Natural Resources	Basic Horticulture
Food System	Irrigation and water management
Crop Production	
Cereals or Basic Grains: Oats, Wheat, Barley, Rye, Triticale, Corn, Sorghum, Millet, Quinoa and Amaranth	Oilseeds: Canola, Rapeseed, Flax, Sunflowers, Corn and Hempseed
Pulse Crops: Peas (all types), field beans, faba beans, lentils, soybeans, peanuts and chickpeas.	Hay and Silage (Forage crop) Production
Vegetable crops or Olericulture: Crops utilized fresh or whole (wholefood crop, no or limited processing, i.e., fresh cut salad); (Lettuce, Cabbage, Carrots, Potatoes, Tomatoes, Herbs, etc.)	Tree Fruit crops: apples, oranges, stone fruit (i.e., peaches, plums, cherries)
Tree Nut crops: Hazlenuts. walnuts, almonds, cashews, pecans	Berry crops: strawberries, blueberries, raspberries
Sugar crops: sugarcane. sugar beets, sorghum	Potatoes varieties and production.
Livestock Production	
Animal husbandry	Ranch
Camel	Yak
Pigs	Sheep
Goats	Poultry
Bees	Dogs
Exotic species	Chicken Growth
Aquaculture	
Fish farm	Shrimp farm
Freshwater prawn farm	Integrated Multi-Trophic Aquaculture
Milk Production (Dairy)	
Dairy goat	Dairy cow
Dairy Sheep	Water Buffalo
Moose milk	Dairy product
Forest Products and Forest management	
Forestry/Silviculture	Agroforestry
Silvopasture	Christmas tree cultivation
Maple syrup	Forestry Growth
Mechanical	
General Farm Machinery	Tillage equipment
Harvesting equipment	Processing equipment
Hay & Silage/Forage equipment	Milking equipment
Hand tools & activities	Stock handling & control equipment
Agricultural buildings	Storage

Agricultural Input Products	
Crop Protection Chemicals	Feed supplements
Chemical based (inorganic) fertilizers	Organic fertilizers
Environmental Science	
Environmental science and regulation	Ecotoxicology
Environmental health issues	Atmosphere and climate
Terrestrial ecosystems	Aquatic ecosystems
Energy and environment	Marine research
Biodiversity	Pharmaceuticals in the environment
Genetically modified organisms	Biotechnology
Risk assessment	Environment society
Theoretical production ecology	horticulture
Breeding	plant fertilization

Board Members

Dr. Bhagawan Bharali (Chief Editor)

Professor & Head, Department of Crop Physiology, Faculty of Agriculture, Assam Agricultural University, Jorhat-785013 (Assam).

Mr. Mukesh Arora (Managing Editor)

M.Tech (Digital Communication), BE (Electronics & Communication), currently serving as Associate Professor in the Department of EE, BIET, Sikar.

Dr. Kusum Gaur (Associate Editor)

Dr. Kusum Gaur working as professor Community Medicine and member of Research Review Board of Sawai Man Singh Medical College, Jaipur (Raj) India.

She has awarded with WHO Fellowship for IEC at Bangkok. She has done management course from NIHF. She has published and present many research paper in India as well as abroad in the field of community medicine and medical education. She has developed Socio-economic Status Scale (Gaur's SES) and Spiritual Health Assessment Scale (SHAS). She is 1st author of a book entitled " Community Medicine: Practical Guide and Logbook.

Research Area: Community Medicine, Biostatistics, Epidemiology, Health and Hospital Management and Spiritual Health

Dr. Darwin H. Pangaribuan

Associate Professor in Department of Agronomy and Horticulture, Faculty of Agriculture, University of Lampung, Indonesia.

Educational background: (Ir.) from Faculty of Agriculture, IPB University, Bogor, Indonesia; (Dipl. Eng) in Land Evaluation from the University of Twente (UT-ITC), Enschede, The Netherlands; (M.Sc) in Crop Production from Wageningen University (WU), The Netherlands. (Ph.D) in Horticulture from University of Queensland (UQ), Brisbane, Australia.

Research Interest: Vegetable Production & Physiology; Biostimulant & Biofertilizers; Organic Farming, Multiple Cropping, Crop Nutrition, Horticulture.

Dr Peni Kistijani Samsuria Mutalib

Working as Research coordinator and HOD in the department of Medical Physics in University of Indonesia.

Professor Jacinta A.Opara

Working full-time and full-ranked Professor and Director, Centre for Health and Environmental Studies at one of the top 10 leading public Universities in Nigeria, the University of Maiduguri-Nigeria founded in 1975.

Dr. Samir B. Salman AL-Badri

Samir Albadri currently works at the University of Baghdad / Department of Agricultural Machines and Equipment. After graduation from the Department of Plant, Soils, and Agricultural Systems, Southern Illinois University Carbondale. The project was 'Hybrid cooling to extend the saleable shelf life of some fruits and vegetables. I worked in many other subject such as Evaporative pad cooling.

Orchid ID: <https://orcid.org/0000-0001-9784-7424>

Publons Profile: <https://publons.com/researcher/1857228/samir-b-albadri>

Dr. Goswami Tridib Kumar

Presently working as a Professor in IIT Kharagpur from year 2007, He Received PhD degree from IIT Kharagpur in the year of 1987.

Prof. Khalil Cherifi

Professor in Department of Biology at Faculty of Sciences, Agadir, Morocco.

Dr. Josiah Chidiebere Okonkwo

PhD Animal Science/ Biotech (DELSU), PGD Biotechnology (Hebrew University of Jerusalem Senior Lecturer, Department of Animal Science and Technology, Faculty of Agriculture, Nau, AWKA.

Prof. Özhan ŞİMŞEK

Agriculture Faculty, Department of Horticulture, Çukurova University, Adana, 01330 Turkey.

Dr. Anka Ozana Čavlović

Working as Professor in the department of Faculty of Forestry, University of Zagreb, Svetošimunska 25, Zagreb.

Dr. Rakesh Singh

Professor in Department of Agricultural Economics, Institute of Agricultural Sciences, Banaras Hindu University, Also Vice President of Indian Society of Agricultural Economics, Mumbai.

Dr. Sunil Wimalawansa

MD, PhD, MBA, DSc, is a former university professor, Professor of Medicine, Chief of Endocrinology, Metabolism & Nutrition, expert in endocrinology; osteoporosis and metabolic bone disease, vitamin D, and nutrition.

Dr. Ajeet singh Nain

Working as Professor in GBPUA&T, Pantnagar-263145, US Nagar, UK, India.

Dr. Salvinder Singh

Presently working as Associate Professor in the Department of Agricultural Biotechnology in Assam Agricultural University, Jorhat, Assam.

Dr. Salvinder received MacKnight Foundation Fellowship for pre-doc training at WSU, USA – January 2000- March 2002 and DBT overseas Associateship for Post-Doc at WSU, USA – April, 2012 to October, 2012.

Dr. V K Joshi

Professor V.K.Joshi is M.Sc., Ph.D. (Microbiology) from Punjab Agricultural University, Ludhiana and Guru Nanak Dev University, Amritsar, respectively with more than 35 years experience in Fruit Fermentation Technology, Indigenous fermented foods, patulin ,biocolour ,Quality Control and Waste Utilization. Presently, heading the dept. of Food Science and Technology in University of Horticulture and Forestry, Nauni-Solan (HP), India.

Dr. Mahendra Singh Pal

Presently working as Professor in the dept. of Agronomy in G. B. Pant University o Agriculture & Technology, Pantnagar-263145 (Uttarakhand).

Dr. Sanjoy Kumar Bordolui

M.Sc. (Ag.), PhD, FSTA, FSIESRP, Assistant Professor, Department of Seed Science and Technology, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia. W.B., India. He received CWSS Young Scientist Award-2016, conferred by Crop and Weed Science Society, received Best Young Faculty Award 2019 conferred by Novel Research Academy, received Innovative Research & Dedicated Teaching Professional Award 2020 conferred by Society of Innovative Educationalist & Scientific Research Professional, Chennai.

Dr. Chiti Agarwal

Dr. Chiti Agarwal works as a postdoctoral associate at the University of Maryland in College Park, Maryland, USA. Her research focuses on fungicide resistance to fungal diseases that affect small fruits such as strawberries. She graduated from North Dakota State University in Fargo, North Dakota, with a B.S. in biotechnology and an M.S. in plant sciences. Dr. Agarwal completed her doctorate in Plant Pathology while working as a research and teaching assistant. During her time as a graduate research assistant, she learned about plant breeding, molecular genetics, quantitative trait locus mapping, genome-wide association analysis, and marker-assisted selection. She wants to engage with researchers from many fields and have a beneficial impact on a larger audience.

DR. Owais Yousuf

Presently working as Assistant professor in the Department of Bioengineering, Integral University-Lucknow, Uttar Pradesh, India.

Dr. Vijay A. Patil

Working as Assistant Research Scientist in Main Rice Research Centre, Navsari Agricultural University, Navsari. Gujarat- 396 450 (India).

Dr. Amit Kumar Maurya

Working as Junior Research Assistant in the Department of Plant Pathology at Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U.P. India.

Prof. Salil Kumar Tewari

Presently working as Professor in College of Agriculture and Joint Director, Agroforestry Research Centre (AFRC) / Program Coordinator in G.B. Pant University of Agric. & Tech., Pantnagar - 263 145, Uttarakhand (INDIA).

Dr. S. K. Jain

Presently working as Officer Incharge of All India Coordinated Sorghum Improvement Project, S. D. Agricultural University, Deesa, Gujarat.

Dr. Deshmukh Amol Jagannath

Presently working as Assistant Professor in Dept. of Plant Pathology, College of Agriculture polytechnic, NAU, Waghai.

Mr. Anil Kumar

Working as Junior Research Officer/Asstt. Prof. in the dept. of Food Science & Technology in Agriculture & Technology, Pantnagar.

Mr. Jiban Shrestha

Scientist (Plant Breeding & Genetics)

Presently working as Scientist (Plant Breeding and Genetics) at National Maize Research Programme (NMRP), Rampur, Chitwan under Nepal Agricultural Research Council (NARC), Singhdarbar Plaza, Kathmandu, Nepal.

Mr. Aklilu Bajigo Madalcho

Working at Jigjiga University, Ethiopia, as lecturer and researcher at the College of Dry land Agriculture, department of Natural Resources Management.

Mr. Isaac Newton ATIVOR

MPhil. in Entomology, from University of Ghana.

He has extensive knowledge in tree fruit orchard pest management to evaluate insecticides and other control strategies such as use of pheromone traps and biological control to manage insect pests of horticultural crops. He has knowledge in agronomy, plant pathology and other areas in Agriculture which I can use to support any research from production to marketing.

Mr. Bimal Bahadur Kunwar

He received his Master Degree in Botany from Central Department of Botany, T.U., Kirtipur, Nepal. Currently working as consultant to prepare CCA-DRR Plan for Hariyo Ban Program/CARE in Nepal/GONESA.

Table of Contents

Volume-10, Issue-1, January 2024

S.No	Title	Page No.
1	Improved Water Quality of Lake Ranisagar using Bioaugmentation Technology Authors: Juan Carlos Verardo, Himanshu Ramesh Lamba  DOI: https://dx.doi.org/10.5281/zenodo.10591064  Digital Identification Number: IJOEAR-JAN-2024-2	01-07
2	Nutrient use Efficiency of different Organic Fertilizer and its Effect on Soil Properties and Yield of <i>Brassica Alboglabra</i> Authors: Ibrahim, I.Z., Sadek, A.R.  DOI: https://dx.doi.org/10.5281/zenodo.10591114  Digital Identification Number: IJOEAR-JAN-2024-9	08-13
3	The Effect of Inoculation Phosphate Solubilizing Bacteria and <i>Rhizobium sp.</i> on Plant Growth and Production of Cowpea (<i>Vigna unguiculata</i>) Authors: Nurul Muddarisna, Yekti Sri Rahayu, Ira Novi Cahyarini  DOI: https://dx.doi.org/10.5281/zenodo.10591146  Digital Identification Number: IJOEAR-JAN-2024-10	14-22

Improved Water Quality of Lake Ranisagar using Bioaugmentation Technology

Juan Carlos Verardo¹, Himanshu Ramesh Lamba^{2*}

BiOWiSH Technologies, 2717 Erie Ave, Cincinnati OH, 45208

*Corresponding Author

Received:- 01 January 2024/ Revised:- 08 January 2024/ Accepted:- 17 January 2024/ Published: 31-01-2024

Copyright © 2024 International Journal of Environmental and Agriculture Research

This is an Open-Access article distributed under the terms of the Creative Commons Attribution

Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted

Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— Bioaugmentation technology achieved rapid bioremediation to significantly improve the water quality of Ranisagar Chaupati Lake over the course of our 8-week study. At the beginning of our study, Ranisagar Chaupati Lake was graded on the Water Quality Index (WQI) as bad water quality and on the Trophic State Index (TSI) as highly eutrophic. Bioaugmentation technology was added to the lake as an all-natural, environmentally safe protocol to improve water quality. This resulted in overall improvement in the WQI from 36.6 (Bad quality) to 57.5 (Moderate quality) and the trophic status of the lake improved from hypereutrophic to eutrophic state.

Keywords— Bioaugmentation, Ranisagar Chaupati, Water Quality, WQI, Eutrophic.

I. INTRODUCTION

The state of Chhattisgarh in central India was created in the year 2000 as a result of the Madhya Pradesh Reorganization Act. It is the 10th largest state in India (by surface) and with a population of over 25 million people it is one of the fastest developing states in India. The state is dotted with numerous lakes, ponds and other water bodies that serve as a major tourist attraction.^[1] Most of the pilgrimage sites in Chhattisgarh are located in the vicinity of a lake or a pond. Human activities in the vicinity of the water bodies have caused deterioration of water quality. The Ranisagar Chaupati – a highly eutrophic lake, located in Rajnandgaon district of the state Chhattisgarh – was suggested by the local municipal corporation as a location to demonstrate Bioaugmentation technology capacity to improve water quality through an all-natural bioremediation protocol.

The Ranisagar Chaupati Lake situated at 21°5'28"N 81°1'22"E in Rajnandgaon district is surrounded by garden which is used for public recreation and is likely to receive a higher nutrient load than any other lake situated in its vicinity.

The lake has a surface area of 20100 m² (2.01 ha) and an average initial water depth of 1.3 m. Stagnant bodies of water such as lakes or ponds are typically more affected by surrounding human activities than flowing surface water (like streams or rivers).

Ranisagar Chaupati is used for public recreation, but the pond has been receiving discharges of domestic sewage and grey waters from an open canal which collects from surrounding settlements (nallah) for over a decade. The main objective behind the project was to improve the overall water quality of Ranisagar Chaupati Lake through the implementation of an all-natural, environmentally safe protocol using Bioaugmentation Technology.

II. MATERIALS AND METHODS

The bioaugmentation technology used for this study is a proprietary composite biocatalyst that enhances a broad range of hydrolytic, oxidative, and reductive biochemical reactions. It contains a novel consortium of metabolically cooperative microorganisms with endogenous and exogenous enzymes, and small molecule co-factors which support both biocatalytic and metabolic activity. They are composed of all-natural materials and non-genetically modified.

Ranisagar Chaupati is a small and shallow lake which continuously received inflow of domestic sewage and grey waters from the nearby households over the past decade. An eight-week bioremediation protocol using Bioaugmentation technology was implemented starting December 28th, 2015. Before starting the study, all of the pond's inlets and outlets were closed and

floating aquatic plants and debris were removed. Samples collected on December 13th, 2015 revealed the following starting water quality:

TABLE 1
DATA FOR INITIAL SAMPLING

Parameter	Unit	Lake condition before bioremediation with BIOWISH* Aqua*	Bathing Water Standards as per MoEf**
PH	mg/l	9.3	6.5-8.5
DO	mg/l	3.6	5 or more
COD	mg/l	88.8	<10
BOD	mg/l	34.7	3 or less
TSS	mg/l	58.5	<10

Table explaining initial sampling

***Results for samples drawn on December 13th, 2015**

****Standards for Bathing water as per MoEF-Ministry of Environment, Forest and Climate Change (Government of India)**

Dosing: The lake was arbitrarily divided into 11 sections to facilitate dosing. An initial shock dose of 56 kg was sprayed on December 28. Lower maintenance doses of 14 kg (aiming to maintain a 0.5 ppm activity for the calculated water volume) were added once a week. The Bioaugmentation technology used for this study is a water-soluble powder and ships ready to dose. For each pond section, the required amount of product was dissolved in pond water and sprayed on the surface using a portable pump. The entire dosing event for the 2-ha pond took no longer than two hours each week.




FIGURE 1: Lake Ranisagar divided into 11 sections

Sampling: CSIR – NEERI: The National Environmental Engineering Research Institute (NEERI) established in Nagpur in 1958, was created and funded by the Government of India (GoI) reporting to the Ministry of Science and Technology (India) of Central Government. NEERI is a pioneer laboratory in the field of environmental science and engineering and part of Council of Scientific and Industrial Research (CSIR) having five zonal laboratories at Mumbai, Kolkata, Hyderabad, Chennai, and Delhi.^[2] NEERI played a major role in auditing the entire project. Since the bioremediation project for Ranisagar Chaupati lake

was approved by the State Government of Chhattisgarh, it was important that the sampling, chain of custody and water quality analysis be done by a reputable GoI certified lab.

TABLE 2
SUMMARY OF DOSING EVENT

Date	Description
December 28th	1st dosing (Shock Dose)
January 7th	2nd dosing - Maintenance Dose
January 14th	3rd dosing Maintenance Dose
January 21st	4th dosing- Maintenance Dose
February 4th	5th dosing Maintenance Dose
February 11th	6th dosing Maintenance Dose
February 18th	7th dosing - Maintenance Dose
February 25th	8th dosing - Maintenance Dose

NEERI's sampling protocol included 22 sampling locations distributed across the pond's area. Grab samples were collected from these locations on the dates shown in the table below. Surface water samples were collected directly without any filtration in acid-rinsed polypropylene bottles, depth water samples were collected approximately 1m from the surface using depth sampler and transferred to clean polypropylene bottles and sludge samples from the bottom of the lake were collected using a van-veen sediment sampler and then transferred into zip-lock polyethylene bags. Similarly, samples for phytoplankton and zooplankton were collected separately and preserved immediately on-site, while samples for chlorophyll a estimation were collected in acid-rinsed amber colored polypropylene bottles and kept away from direct sunlight until analyzed. Separate samples for fecal coliform analysis were collected in sterile polypropylene bottles. Sample pH, temperature and dissolved oxygen were immediately determined on-site, while the remaining samples were shipped for analysis at CSIR-NEERI at Nagpur after adequate preservation of the collected samples. Amongst the 5 sampling events conducted, 2 sampling events (2nd and 4th monitoring) were conducted to analyze 13 major parameters (pH, temperature, DO, COD, BOD, ammonia, nitrate, TKN, phosphate, suspended solids, total dissolved solids, fecal coliform and chlorophyll a) and 3 sampling events (1st, 3rd and 5th monitoring) were conducted to analyze 33 parameters for complete analysis of the lake including water characteristics, sediment characteristics, aquatic diversity and physio-chemical parameters inclusive of above mentioned 13 parameters.




FIGURE 2: Schematic of sampling location in Ranisagar Lake

TABLE 3
SAMPLING DATES

Date	Description
December 13th	1st sampling before dosing (lesser parameters)
January 13th	2nd sampling (full parameters)
January 27th	3rd sampling (lesser parameters)
February 10th	4th sampling (lesser parameters)
February 24th	5th sampling (full parameters)

III. RESULTS AND DISCUSSION

The graphs below show the 22 surface water samples for each sampling event plotted in a “box graph” style. The top and bottom lines for the box represent upper and lower quartiles, while the center line shows the mean (arithmetic average for the 22 samples). The bars extending from the boxes show: Quartile $\pm 1.5 * (\text{Inter Quartile Range})$.




FIGURE 3: Graph 1 Five-point summaries of data collected and Graph 2 Average values of 22 sampling points for BOD




FIGURE 4: Graph 3 Three-point summaries of data collected and Graph 4 Average values of 13 sampling points for % organic fraction

The fluctuations in BOD can be attributed to the rapid degradation of organic matter in the system. This increased biological activity in the sediment may have resulted in changing the release of organic components which contributed to varying BOD readings.

Sludge in sewage treatment lagoons are likely to contain over 50% organic fraction. Old material which is allowed to mineralize for years may have 10 to 15% organic fractions. Chaupati pond sludge was found to have an initial 8.7% organic fraction, with very high levels of Fe and Mn – indicating iron rich sandy soils and a very advanced stage of mineralization, a very poor indication for bioremediation. Despite this, rapid action could be seen on the sludge by week 4. A 45% drop in sludge organic fraction was seen 4 weeks into the program. Increased biological activity is also reflected in the TSS spike. Achieving lower BOD and sludge organic fraction values in 8 weeks shows how rapidly Bioaugmentation technology can catalyze biological action in water systems.




FIGURE 5: Graph 5 Five-point summaries of data collected and Graph 6 Average values of 22 sampling points

Bioaugmentation technology used for this study contains a consortium of microorganisms which can facilitate diverse nitrogen management metabolic pathway in water bodies. The microbial consortium in the bioaugmentation technology is able to promote ammonia (NH_3) oxidation directly to gaseous forms of nitrogen through Heterotrophic Nitrification and Denitrification (HND).^[3] This results in lower ammonia nitrogen concentrations while avoiding peak formations of nitrite and nitrate (which may be toxic to aquatic life). The different oxidative/reductive pathways in HND by-pass the use of dissolved oxygen as an electron acceptor in aerobic nitrification, resulting in higher DO levels for the pond (as seen on Graph 14).




FIGURE 6: Graph 7 Five-point summaries of data collected and Graph 8 Average values of 22 sampling points for NO_3



FIGURE 7: Graph 9 Five-point summaries of data collected and Graph 10 Average values of 22 sampling points for TKN




FIGURE 8: Graph 11 Five-point summaries of data collected and Graph 12 Average values of 22 sampling points for NH_3




FIGURE 9: Graph 13 Five-point summaries of data collected and Graph 14 Average values of 22 sampling points for DO

We see an increase in phosphate concentrations which may be explained by the increased release of nutrients from the lake sediment via aggressive decomposition of organic fraction. A sharp drop in phosphate concentration could be observed after the 5th monitoring, indicating that the slower rate of organic degradation in the sludge allows for more effective phosphate settling.

During the study, the pH of the lake decreased consistently from an average value of 9.3 to 8.8. Decrease in pH indicates improved biological activity^[4]. Bioaugmentation also showed great effect on restoring the pH values in the lake water.




FIGURE 10: Graph 15 Five-point summaries of data collected and Graph 16 Average values of 22 sampling points for $\text{PO}_4\text{-P}$




FIGURE 11: Graph 17 Five-point summaries of data collected and Graph 18 Average values of 22 sampling points for pH

3.1 Water Quality Index (WQI)

WQI, developed by National Sanitation Foundation (NSF-WQI), was used to quantify the change in the quality of water in response Bioaugmentation. NSF-WQI categorizes water quality into five categories: 0-25 Very Bad, 25-50 Bad, 50-70 Moderate, 70-90 Good and 90-100 Excellent. In this method, nine different physio-chemical parameters are used to determine the WQI (dissolved oxygen, fecal coliform, pH, BOD, temperature, total phosphate, nitrate, turbidity, and total solids).^[5]

The overall changes in the WQI of the lake during 1st monitoring to 5th monitoring is depicted as follows:

TABLE 4
CHANGES IN WQI OF LAKE DURING 1ST MONITORING TO 5TH MONITORING

Sampling	1 st	2 nd	3 rd	4 th	5 th
NSF-WQI	36.6	46.6	55.1	56.1	57.5
Water Quality	Bad	Bad	Moderate	Moderate	Moderate

3.2 Trophic State Index (TSI)

The Trophic State Index (TSI) classifies the trophic status of a lake or any water body using a measurable value. The most commonly used numerical classification method of the trophic status of a lake was that developed by Robert Carlson in the year 1977 and is known as Carson's Trophic State Index (TSI). Carson's TSI uses three dependent variables which have direct correlation with the trophic status of the lake, namely, algal biomass, transparency of the lake water (which enables light penetration), and the nutrient concentration to classify the trophic status of the lake into hyper oligotrophic, oligotrophic, mesotrophic, eutrophic, and hypereutrophic conditions.^[6]

TABLE 5
CLASSIFICATION OF LAKES BASED ON TSI VALUES^[6]

TN values	Trophic Status
<30	Hyperoligotrophic
30—40	Oligotrophic
40-50	Mesotrophic
50-70	Eutrophic
>70	Hypereutrophic

Based on the TSI values, Ranisagar Chaupati Lake was classified as hypereutrophic before starting the Bioaugmentation. The water quality improved significantly after Bioaugmentation and resulted in the trophic status of the lake to shift from hypereutrophic to eutrophic state in 8 weeks.

IV. CONCLUSION

Bioaugmentation in Ranisagar Chaupati Lake helped to remarkably improve the water quality. We saw an overall improvement in the water quality index (WQI) from 36.6 (Bad quality) to 57.5 (Moderate quality) and the trophic status of the lake improved from hypereutrophic to eutrophic state. The concentration of chlorophyll a, an algal pigment that imparts green color to the water^[7], decreased by 40%. Decrease in concentration of Chlorophyll a was due to significant change in the population density (size) and dynamics of phytoplankton (algal) species. 21% reduction was observed in the phytoplankton population after Bioaugmentation. The reduction in population of phytoplankton was reflected in reduction in pH of the lake water from an average value of 9.3 to 8.8. Very high pH in lake bodies results due to high photosynthetic activity^[8]. Bioaugmentation also caused significant reduction in the concentrations of organic ammonia (represented as TKN), inorganic ammonia and nitrate by 65%, 40% and 79%, respectively. The organic carbon concentration of lake water, as represented in terms of BOD concentrations also showed an average reduction of 29% within two months of Bioaugmentation.

Similarly, the lake sediment also showed a 72% reduction in organic fraction during this period. Improvement in the water quality is also reflected by an overall decrease in the amount of pollution indicating algal species in lake water as represented by Palmer's Pollution Index (PPI). The PPI of the lake decreased by 47% indicating improvement of lake water quality in terms of reduced nutrient concentration.

TABLE 6
EFFECT OF BIOAUGMENTATION ON SEVERAL PARAMETERS BEFORE AND AFTER DOSING

Parameters	Before Dosing (13th Dec 2015)	After Dosing (24th Feb 2016)	Observation	Remarks
pH	9.3	8.8	Decreased by 5%	Improved
Dissolved Oxygen	3.6	8.4	Increased by 133%	Improved
Chlorophyll a	0.30	0.18	Decreased by 40%	Improved
TKN (as N)	98.7	34.5	Decreased by 65%	Improved
Ammonia (as N)	9.8	5.9	Decreased by 40%	Improved
Nitrate (as NO ₃)	2.4	0.5	Decreased by 79%	Improved
Phosphate (as P)	1.0	0.6	Decreased by 40%	Improved
BOD (3d, 27°C)	35	25	Decreased by 29%	Improved
Sediment Organics (%)	8.7	2.4	Decreased by 72%	Improved
Phytoplankton (Density/L)	7.7×10^6	6.1×10^6	Decreased by 21 %	Improved
Palmer's Pollution Index (PPI)	19	10	Decreased by 47%	Improved
Zoopankton (Density/L)	81333	81	Decreased by 99.9%	Improved

ACKNOWLEDGEMENT

We would like to thank Bridgedots Techservices Pvt. Ltd., Effioa and CSIR-NEERI for their help and support in carrying out the field activities and data compilation for this study.

REFERENCES

- [1] "Chhattisgarh," Wikipedia, Oct. 25, 2020. <https://en.wikipedia.org/wiki/Chhattisgarh>
- [2] "National Environmental Engineering Research Institute," Wikipedia, Dec. 17, 2023. https://en.wikipedia.org/wiki/National_Environmental_Engineering_Research_Institute (accessed Dec. 18, 2023).
- [3] J. K. Kim, K. J. Park, K. S. Cho, S.-W. Nam, T.-J. Park, and R. Bajpai, "Aerobic nitrification–denitrification by heterotrophic *Bacillus* strains," *Bioresource Technology*, vol. 96, no. 17, pp. 1897–1906, Nov. 2005, doi: <https://doi.org/10.1016/j.biortech.2005.01.040>.
- [4] B. M. Saalidong, S. A. Aram, S. Otu, and P. O. Lartey, "Examining the dynamics of the relationship between water pH and other water quality parameters in ground and surface water systems," *PLOS ONE*, vol. 17, no. 1, p. e0262117, Jan. 2022, doi: <https://doi.org/10.1371/journal.pone.0262117>.
- [5] R. 10 US EPA, "Freshwater Quality," US EPA, May 14, 2013. <https://www.epa.gov/salish-sea/freshwater-quality>
- [6] R. E. Carlson, "A trophic state index for lakes1," *Limnology and Oceanography*, vol. 22, no. 2, pp. 361–369, Mar. 1977, doi: <https://doi.org/10.4319/lo.1977.22.2.0361>.
- [7] National Geographic, "Chlorophyll | National Geographic Society," [education.nationalgeographic.org](https://education.nationalgeographic.org/resource/chlorophyll/), Jul. 15, 2022.
- [8] S. Zerveas, M. S. Mente, D. Tsakiri, and K. Kotzabasis, "Microalgal photosynthesis induces alkalization of aquatic environment as a result of H⁺ uptake independently from CO₂ concentration – New perspectives for environmental applications," *Journal of Environmental Management*, vol. 289, p. 112546, Jul. 2021, doi: <https://doi.org/10.1016/j.jenvman.2021.112546>.

Nutrient use Efficiency of different Organic Fertilizer and its Effect on Soil Properties and Yield of *Brassica Alboglabra*

Ibrahim, I.Z.^{1*}, Sadek, A.R.²

Organic Farming Program, Soil Science, Fertilizer and Water Research Centre, Malaysian Agricultural Research and Development Institute (MARDI), Serdang, Selangor, Malaysia

*Corresponding Author

Received:- 02 January 2024/ Revised:- 10 January 2024/ Accepted:- 19 January 2024/ Published: 31-01-2024

Copyright @ 2024 International Journal of Environmental and Agriculture Research

This is an Open-Access article distributed under the terms of the Creative Commons Attribution

Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted

Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— In organic farming, homemade organic fertilizer such as compost or enriched organic material had become one of crucial factors for nutrient management. Other than the limited source of N for organic fertilizer as the key factor for growth-limiting nutrients in organic farming, the cost of fertilization is also affected by different organic fertilizer application rates. Therefore, the objectives of this study were (i) to determine the effect of these organic fertilizers on the soil properties (chemical and biological properties) and (ii) to evaluate the efficiency and growth performance of Kailan (*Brassica alboglabra*) as influence by the application of different organic fertilizer in the field. A study was conducted at an integrated organic farming research area in the Malaysian Agricultural Research and Development Institute (MARDI) at Serdang, Selangor for two consecutive cycles. The experimental treatments were focused on the different rates of organic fertilizer inputs (3-33 t/ha from organic sources) with soil alone as a control. The treatments were applied 14 days after transplanting using *Brassica alboglabra* as a test crop. The study result found that a BIOC at 2 tonne kg/ha give significant different in nutrient use efficiency (NUE), however NC at 33 tonne kg/ha rate of organic input had improved most of soil properties NF showed a great yield as well as their growth performances. Study findings have shown that the range of application rate of organic input depends on applicability of the organic fertilizer whether it can promotes nutrient uptake in plants and or can be scaled up to farm level to sustain organic system productivity.

Keywords— Organic Fertilizer, Soil Properties, Yield, Nutrient Use, Organic Farming, Compost.

I. INTRODUCTION

In organic vegetable cultivation, inorganic fertilizer is not allowed with exception of naturally occurring mineral sources. The substitutes for inorganic fertilizer are mostly in organic nutrient sources that can be grouped into such as animal sources, plant sources, green manures and compost (Aini *et al.*, 2005) which can serve as alternative practice in organic crop production. To improve soil fertility, application of organic fertilizer has become an important practical measure. Organic fertilizers can enhance soil microbial activity by promoting the utilization of nutrients and restoration of microbial community structure and function in the rhizosphere soils. They can increase the mass and diversity of the soil microbiota, improve soil fertility, and stimulate soil enzyme activity, ultimately driving the carbon and nitrogen cycles (Huang *et al.*, 2023). Additionally, organic fertilizers can introduce beneficial microorganisms into the soil, increase microbial diversity, and influence the structure and function of nitrifying and denitrifying microorganisms. Therefore, organic fertilizers play a significant role in enhancing soil microbial activity, ultimately contributing to improved soil health and agricultural productivity (Wang *et al.*, 2013; Pu *et al.*, 2022; Sivojiene *et al.*, 2021).

Organic fertilizer is able to enhance soil microbial activity (Ren *et al.*, 1996), increased microbial biomass (Suresh *et al.*, 2004) and also improved soil structure (Bin, 1983; Dauda *et al.*, 2008). Though high amounts of manure or composts are required initially at organic farm, the quantity of subsequent application may be reduce as soil physical, chemical and biological improve with time (Aini *et al.*, 2005). Organic fertilizers can have varying levels of nutrient use efficiency depending on the type of fertilizer and the farming system. Organic farming systems are highly dependent on organic fertilizers, such as manure and compost, which can provide a range of nutrients to the soil (Chmelikova *et al.* 2021). However, the nutrient content of organic fertilizers can be highly variable, and their release rates can be slower than synthetic fertilizers, which can impact nutrient

uptake efficiency (Barlog, 2023). Additionally, organic fertilizers can contribute to soil organic matter, which can improve soil structure and water-holding capacity, ultimately contributing to improved nutrient uptake efficiency (Xingxua et al., 2023).

Overall, the nutrient use efficiency of organic fertilizers can be improved through careful management practices, such as optimizing fertilizer application rates, timing, and placement, and through the use of complementary nutrient sources, such as cover crops and crop rotations (Oluwadunsin et al., 2022). Invariably, many cultivated soils of the world are scarce in one or more of the essential nutrients required supporting vigorous plants. Estimates of overall efficiency of applied fertilizer have been reported to be around or lesser than 50% for N, less than 10% for P, and about 40% for K (Baligar *et al.*, 2001). Plants that are capable in absorption and utilization of nutrients prominently enhance the efficiency of applied fertilizers, decreasing cost of inputs, and avoiding losses of nutrients to ecosystems. To increase fertilizer use efficiency (FUE) and to reduce its negative influence on the atmosphere has been an important point in the world for a long time. It can be particularly affected by fertilizer management as well as by soil and irrigation Management. There is increased demand for fertilizer nutrients to meet the global demand for food, However there are inadequate fertilizer resources available and rising public concern related to nutrient use side effects. This has led to calls for Nutrient use efficiency (NUE) to be improved, but not at the expense of decreased crop productivity. NUE depends on the plant's capacity to take up nutrients efficiently from the soil, but also depends on inner transport, storage and remobilization of nutrient.

II. MATERIAL AND METHODS

2.1 Field experiment and treatment used:

The experiment was conducted at Integrated Organic Farm, MARDI Serdang. The test crop is Kailan was grown under rainshelter and treated with seven different application of organic fertilizer. There were chicken dung (CD), indigenous microorganism compost (IMO), vermicompost (V), normal compost (NC), rice husk charcoal compost (BIOC), nature farming (NF), depleted oil bleaching earth compost (DOBE) and plus untreated soil served as CONTROL. The organic fertilizers were applied according to manual and nutrient requirement for kailan and as shown in Table 1. The following treatments were made in a randomized block design (RCBD) in five replicates.

2.2 Soil sampling and Laboratory Analyses:

Soil at depth 0-20 cm were sampled before organic fertilizers application in composite soil samples and at after harvesting in six replicates for chemical analyses for soil quality. The soils were air dried and ground to pass a 2 mm sieve for analyses. Total nitrogen in soils was analyzed by Micro Kjeldahl digestion followed by distillation and titration with 0.1 N HCL. Organic carbon was determined by Walkey and Black rapid titration method. Available phosphorus in soils was extracted based on the Bray and Kurtz no.2 procedure. The soil acidity was measured on pH meter using 1:2.5 ratio and water. The exchangeable cations were extracted by leaching with 1 N ammonium acetate and their concentration was determined with Inductively Coupled Plasma (ICP) Optical Emission Spectrophotometer. Wet samples of soil were used for microbial population studies of total colony forming unit (cfu-log₁₀) using total plate count on nutrient agar.

2.3 Nutrient use efficiency calculation

Nutrient use efficiency (NUE) may be defined as yield per unit fertilizer input or in terms of recovery of applied fertilizer. Nutrient use efficiency (NUE) is a critically important concept in the evaluation of crop production systems. The equation used as follows;

$$\text{Partial Factor Productivity (PFP)} = Y/F \quad (1)$$

Whereas Y =yield of harvested portion of crop with nutrient applied and F = amount of nutrient applied

2.4 Statistical Analysis

All data were subjected to a one-way analysis of variance (ANOVA). Treatment means were compared using Duncan Multiple Range test at a significance level of $P < 0.05$ using SAS 9.4.

III. RESULTS AND DISCUSSION

Six local made organic fertilizers (NF, CD, NC, VC, IMO, BIOC, DOBE) with its nutrient properties showed in Table 1. Soil properties of chemical characteristic before the application of organic fertilizer to the soil shown in Table 2. At the end of experiment, several properties of soil nutrient status such as in pH, nitrogen content, CEC and all exchangeable cation (K, Na, Ca and Mg) in those treatment increased significantly as showed in Table 3. Treatment with CD (chicken dung) gave most

significant effect in soil nutrient properties. While, the carbon content in all treatment showed no significant different even though there quite high amount of carbon source added in the soil. This might be due to the availability of nutrients especially nitrogen and could be due to improvement of soil water holding capacity.

TABLE 1

SOME CHEMICAL CHARACTERISTIC OF SEVEN LOCAL MADE ORGANIC FERTILIZER USED IN THE EXPERIMENT

Types of fertilizer	pH (H ₂ O)	Organic carbon (%)	Total N (%)	P ₂ O ₅ (%)	K ₂ O (%)	Application tonne/ha
V	6.69	21.32	1.79	1.02	1.75	3
IMO	6.36	20.87	1.983	0.25	0.98	20
DOBE	5.58	4.45	0.44	2.5	0.7	20
NC	6.35	24.49	1.57	0.77	2.83	33.33
CD	5.66	3	1.72	1.82	2.18	33.33
BIOCHAR	6.5	19.2	4.1	3.3	2.7	2
NF	6.03	1.17	0.27	0.25	0.98	13.33

TABLE 2

SOME CHEMICAL CHARACTERISTIC OF THE SOIL BEFORE APPLICATION OF ORGANIC FERTILIZER

Soil parameters	Values
Soil pH (H ₂ O)	6.75
CEC	6.21
Conductivity (µs/cm)	1108
Organic carbon (%)	3.27
Nitrogen (%)	0.118
P (ug/gm)	406.05
Exchangeable K (meq/100gm soil)	101.41
Exchangeable Na (meq/100gm soil)	257.18
Exchangeable Ca (meq/100gm soil)	11.8
Exchangeable Mg (meq/100gm soil)	52.48
Total microbial count (log ₁₀ CFU)	5.792

The total microbial count in Table 3 does not reflect any variation and had no significant different effect from any organic fertilizer type that had been used compared to control. Generally, the plate culture method normally can only separate 0.1-1.0% of soil microbes and cannot reflect the original status of soil microbial diversity (Cai and Liao, 2002; Vigdis and Lisc, 2002; Luo *et al.* 2003). While, according to vanZwieten (2006), it explained as variable effect of a given amendment on different organism may change the composition of the microbial community without changing the total population or activities.

Management in the use of fertilizer even organic fertilizer can be increase through few recommendation such as use it at right time which is the interrelated to site specific nutrient management (SSNM). Greater synchronization between crop demand and nutrient supply is necessary to improve nutrient use efficiency, especially for N (Giller *et al.*, 2004). Split applications of N during the growing season, rather than a single, large application prior to planting, are known to be effective in increasing N use efficiency. Besides that, integrated nutrient management (INM) involves optimum use of indigenous nutrient components i.e. crop residues, organic manure, biological N fixation as well as chemical fertilizer and their balancing interactions to rises N and P recovery. The appropriate understanding and exploitation of these positive interactions among the plant nutrient is keys for increasing returns to the farmers in terms of yield as well as soil quality and NUE of applied N (Aulakh and Malhi, 2004).

TABLE 3
EFFECT OF DIFFERENT ORGANIC FERTILIZERS ON SOIL PROPERTIES (CHEMICAL AND BIOLOGICAL) AFTER HARVEST PERIOD

Treatments	pH H ₂ O	Conductivity μ S	%		SOIL - P uq/qm of soil	Total microbial count (Log ₁₀ CFU)
			Nitrogen	Carbon		
NF	7.06 b	397.7 c	0.09 b	2.69 a	334.34 ab	6.275 a
BIOC	7.03 bc	861.2 b	0.09 b	2.85 a	341.56 ab	6.270 a
NC	7.46 a	1615.8 a	0.14a	2.77 a	244.70 b	6.340 a
CONTROL	7.06 b	1152.1 b	0.10 b	2.59 a	315.18 ab	6.363 a
IMO	7.13 b	720.0 bc	0.09 b	2.40 a	292.22 ab	6.318 a
V	7.16 b	804.0 bc	0.11 ab	2.59 a	296.10 ab	6.248 a
CD	7.13 b	909.2 b	0.12 ab	2.46 a	363.24 a	6.521 a
DOBE	6.78 c	963.0 b	0.11 ab	2.68 a	305.80 ab	6.027 a
Treatments	meq/100q soil					
	CEC	Exch K	Exch Ca	Exch Na	Exch Mg	
N F	11.14 ab	65.32 b	163.64 a	5.02 c	34.03 c	
BIOC	10.96 ab	106.48 b	150.97 a	8.61 bc	34.46 c	
NC	14.37 a	238.37 a	145.07 a	18.02a	47.84 ab	
CONTROL	8.19 b	133.56 b	193.53 a	10.03 bc	43.40 ab	
IMO	12.15 ab	112.79 b	142.79 a	6.60 c	31.97 b	
V	11.56 ab	119.79 b	174.99 a	8.75 bc	39.41 ab	
CD	14.09 a	137.55 b	226.56 a	14.27 ab	56.29 a	
DOBE	9.72 ab	82.45 b	238.54 a	8.17 bc	43.88 ab	

Values are given as a mean of five replicates. Means with the same letter in a column are not significantly different ($P < 0.05$) as determined by Duncan Multiple Range Test




FIGURE 1: Graph on yield of kailan and fertilizer use efficiency (NUE)

The projected yield of kailan as shown in Figure 1 is estimated as medium-low production per hectare of season. The total average yield is about 8490 kg/ha below the targeted medium production 9000 kg/ha. Yield of kailan per plot were higher in treatment with NF (Nature Farming) with estimation of up to 12000 kg per hectare. In term of nutrient use efficiency (NUE) with partial factor of productivity (kg crop yield per kg nutrient applied) as described by Mosier *et al* 2004, it showed treatment with BIOC (biochar compost) and V (vermicompost) gave higher value index compared to other treatment.

The use of organic fertilizers can have several positive effects on soil properties as showed in Table 3. Organic fertilizers can improve soil structure, texture, and aeration, leading to better water retention and healthy root growth (Joaqim et al., 2023; Zhou et al, 2022). They also contribute to the long-term availability of nutrients, add beneficial microorganisms to the soil, and increase soil organic carbon and other nutrients (Sissay, 2019). Additionally, organic fertilizers are environmentally friendly, provide micronutrients, and are less likely to cause burning to plants. Overall, the application of organic fertilizers can lead to improved soil quality, increased crop yields, and sustainable agricultural.

IV. CONCLUSION

The use of different organic fertilizers in the cultivation of *Brassica alboglabra* can significantly influence nutrient use efficiency, soil properties, and crop yield. The effectiveness of these fertilizers depends on their composition and the soil's existing conditions. Organic fertilizers typically improve soil health by enhancing its physical structure, increasing microbial activity, and supplying essential nutrients. This leads to better nutrient availability and uptake by plants, often resulting in improved growth and yield of *Brassica alboglabra*. Moreover, the use of organic fertilizers is also beneficial for long-term soil fertility and environmental sustainability. However, specific outcomes can vary based on the type of organic fertilizer used, application rates, and the particular soil and environmental conditions. Application with organic fertilizer to the soil had increased most of its nutrients properties. Leafy vegetables such as for this experiment, Kailan (*Brassica alboglabra*) can be grown better in soil amended with organic fertilizer but the application rate and availability of all minerals should be considered. Furthermore, organic leafy vegetables are expected to be a quality product for human to consume and maybe profitable to producer and even environmental than those from conventional production system. Further studies are needed to determine optimal rates of organic fertilizer for proper growth and production of organic leafy vegetables.

ACKNOWLEDGEMENTS

The authors would like to thank the Director General of MARDI and the Director of Soil Science, Fertilizer and Water Research Centre for their support. We thank also to Dzulkifli Mat Lajis, Muhammad Hazwan Husaimi and Muhammad Noor Azmie Abdul Jalil Salleh for their assistance throughout the field experiment and laboratory analyses. This project was funded under RMK10-150 by Government of Malaysia.

REFERENCES

- [1] Aini, Z., Sivapragasam, A., Vimala, P., and Mohamad Roff, M.N. (2005). Organic vegetable cultivation in Malaysia. Malaysian Agricultural Research and Development Institute. Pp 47-67.
- [2] Aulakh and Malhi. (2004). Fertilizer N use efficiency as influenced by interactions with other nutrients, In: Agriculture and the Nitrogen Cycle: Assessing the Impacts of Fertilizer Use on Food Production and the Environment (Editors: A.R. Mosier, J.K. Syers and J.R. Freney); 181-191. Island Press, Washington, USA
- [3] Baligar, V.C., Fageria, N.K. and He, Z.L. (2001). Nutrient use efficiency in plants. *Communications in Soil Science and Plant Analysis*. 32(7&8), 921–950.
- [4] Huang, B., Chen, Y., Pei, Z., Jiang, L., Zhang, Y., Wang, J., & Wang, J. (2023). Application of microbial organic fertilizers promotes the utilization of nutrients and restoration of microbial community structure and function in rhizosphere soils after dazomet fumigation. *Frontiers in Microbiology*, 13, 1122611. <https://doi.org/10.3389/fmicb.2022.1122611>
- [5] Barlóg P. (2023). Improving Fertilizer Use Efficiency-Methods and Strategies for the Future. *Plants* (Basel). 12(20):3658. doi: 10.3390/plants12203658. PMID: 37896121; PMCID: PMC10610152.
- [6] Bin, J. 1983. Utilization of green maize for raising soil fertility in China. *Soil Sci.*, 135:65-9.
- [7] Cai. Y.F and Liao, Z.W. 2002. Advance in research methods for soil microbial ecology. *Soil and Environment II*, 167-171.
- [8] Chmelíková, L., Schmid, H., Anke, S. and Hulsbergen, K.J. 2021 Nitrogen-use efficiency of organic and conventional arable and dairy farming systems in Germany. *Nutr Cycl Agroecosyst* **119**, 337–354 (2021). <https://doi.org/10.1007/s10705-021-10126-9>
- [9] Dauda, S.N., F.Ajayi and E.E.Ndor. 2008. Growth and yield of water melon (*Citrullus lanatus*) as affected by poultry manure application. *J.Agric. Soc Sci.*, 4:121-4.
- [10] Giller, K. E., Chalk, P., Dobermann, A., Hammond, L., Heffer, P., Ladha, J. K., Nyamudeza, P., Maene, L., Ssali, H., & Freney, J. (2004). Emerging Technologies to Increase the Efficiency of Use of Fertilizer Nitrogen. In A. R. Mosier, J. K. Syers, & J. R. Freney (Eds.), *Agriculture and the Nitrogen Cycle: Assessing the Impacts of Fertilizer Use on Food Production and the Environment* :35-51. (Scope; No. 65).
- [11] Joachim G.C., Deru, J. B., Ron D. G., Lijbert B. and Nick V. E.(2023). Effects of organic and inorganic fertilizers on soil properties related to the regeneration of ecosystem services in peat grasslands. *Applied Soil Ecology*. Volume 187-104838.ISSN 0929-1393, <https://doi.org/10.1016/j.apsoil.2023.104838>.
- [12] Lazcano C, Zhu-Barker X, Decock C. (2021). Effects of Organic Fertilizers on the Soil Microorganisms Responsible for N₂O Emissions: A Review. *Microorganisms*. 1;9(5):983. doi: 10.3390/microorganisms 9050983.

- [13] Luo, H.F., Qi, H.Y and Xue,C. 2003. A preliminary application of PCR-DGGE to study microbial diversity in soil. *Acta Ecological Sinica* 23, 1570-1575.
- [14] Mosier, A.R., J.K. Syers and J.R. Freney. 2004. Agriculture and the nitrogen cycle. Assessing the impacts of fertilizer use on food production and environment. Scope 65. *Island Press, London*.
- [15] Oluwadunsin O., Nanthi B. and Greg H. 2022. A comprehensive review on enhancing nutrient use efficiency and productivity of broadacre (arable) crops with the combined utilization of compost and fertilizers, *Journal of Environmental Management*. Volume 317-115395, ISSN 0301-4797, <https://doi.org/10.1016/j.jenvman.2022.115395>.
- [16] Pu, R., Wang, P., Guo, L., Li, M., Cui, X., Wang, C., Liu, Y., & Yang, Y. (2022). The remediation effects of microbial organic fertilizer on soil microorganisms after chloropicrin fumigation. *Ecotoxicology and Environmental Safety*, 231, 113188. <https://doi.org/10.1016/j.ecoenv.2022.113188>
- [17] Sisay Assefa .(2019). The Principal Role of Organic Fertilizer on Soil Properties and Agricultural Productivity -A Review. *Agri Res& Tech: Open Access J*. 2019; 22(2): 556192. DOI: 10.19080/ARTOAJ.2019.22.556192
- [18] Sivojiene, D., Kacergius, A., Baksiene, E., Maseviciene, A. and Zickiene, L. (2021) The Influence of Organic Fertilizers on the Abundance of Soil Microorganism Communities, Agrochemical Indicators, and Yield in East Lithuanian Light Soils. *Plants (Basel)*. 2;10(12):2648. doi: 10.3390/plants10122648. PMID: 34961119; PMCID: PMC8703430.
- [19] Suresh, K.D., G. Sneh, K.K.Krishn and C.M.Mool, 2004. Microbial biomass carbon dan microbial activities of soils receiving chemical fertilizer and organic amendments. *Archives Agron. Soil Sci.*, 50:641-7
- [20] vanZwieten, L. 2006. Impact of agriculture inputs on soil organism. *Aust. J. Soil Res*. 3:1-15.
- [21] Vigdis, T and Lisc, O. 2002. Microbial diversity and fuction in soil: from gene to ecosystem. *Current Opinion in Microbiology*. 5: 240-245.
- [22] Xingjuan Z., Gerard H. R., Minggang X., Zejiang C., N. S., Yinghua. D. and Wim D.V. (2023) Long-term impacts of mineral and organic fertilizer inputs on nitrogen use efficiency for different cropping systems and site conditions in Southern China. *European Journal of Agronomy*, Volume 146(2023)-126797, ISSN 1161-0301 <https://doi.org/10.1016/j.eja.2023.126797>.
- [23] Wang, Y., Li, Q. and Li, C. (2023). Organic fertilizer has a greater effect on soil microbial community structure and carbon and nitrogen mineralization than planting pattern in rainfed farmland of the Loess Plateau. *Frontiers in Environmental Science*, 11, 1232527. <https://doi.org/10.3389/fenvs.2023.1232527>
- [24] Zhang X., Li J., Shao L., Qin F., Yang J., Gu H., Zhai P. and Pan X. (2023). Effects of organic fertilizers on yield, soil physico-chemical property, soil microbial community diversity and structure of *Brassica rapa* var. *Chinensis*. *Frontiers in Microbiology* (14) <https://www.frontiersin.org/articles/10.3389/fmicb.2023.1132853> DOI=10.3389/fmicb.2023.1132853
- [25] Zhou, Z., Zhang, S., Jiang, N., Xiu, W., Zhao, J., & Yang, D. (2022). Effects of organic fertilizer incorporation practices on crops yield, soil quality, and soil fauna feeding activity in the wheat-maize rotation system. *Frontiers in Environmental Science*, 10, 1058071. <https://doi.org/10.3389/fenvs.2022.1058071>.

The Effect of Inoculation Phosphate Solubilizing Bacteria and *Rhizobium sp.* on Plant Growth and Production of Cowpea (*Vigna unguiculata*)

Nurul Muddarisna¹, Yekti Sri Rahayu^{2*}, Ira Novi Cahyarini³

Agrotechnology Department, Wisnuwardhana University, Danau Sentani Street 99 Malang City, East Java, Indonesia

*Corresponding Author

Received:- 04 January 2024/ Revised:- 13 January 2024/ Accepted:- 23 January 2024/ Published: 31-01-2024

Copyright @ 2024 International Journal of Environmental and Agriculture Research

This is an Open-Access article distributed under the terms of the Creative Commons Attribution

Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted

Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— The constraints of Alfisol or Mediterranean soils pose unique management challenges for maintaining soil fertility and agricultural productivity. The aim of the research was to study the effect of couple inoculation *Rhizobium* and Phosphat Solubilizing Bacteria (PSB) on plant growth and production of Cowpea in Alfisol or Mediterranean soils. The experiment used a Randomized Block Design with treatment as combination concentration of PSB and *Rhizobium sp.* The treatment consist of : control; PSB of 10 ml/L and *Rhizobium sp* 10 ml/kg seed; PSB of 10 ml/L and *Rhizobium sp* 20 ml/kg seed; PSB of 10 ml/L and *Rhizobium sp* 30 ml/kg seed; PSB of 20 ml/L and *Rhizobium sp* 10 ml/kg seed; PSB of 20 ml/L and *Rhizobium sp* 20 ml/kg seed; PSB of 20 ml/L and *Rhizobium sp* 30 ml/kg seed; PSB of 30 ml/L and *Rhizobium sp* 10 ml/kg seed; PSB of 30 ml/L and *Rhizobium sp* 20 ml/kg seed; PSB of 30 ml/L and *Rhizobium sp* 30 ml/kg of seed. Each treatment was repeated with three repetitions. Data analysis used analysis of variance at the 5% level, followed by LSD test at the 5% level. The results showed that couple inoculation PSB and *Rhizobium sp.* increase significantly N and P soil, plant growth of Cowpea i.e the plant height, number of root nodules, fresh weight, shoot dry weight and total dry weight, the average 100 seed weight, population of PSB in soils and *Rhizobium* in root cowpea compared by control.

Keywords— Alfisol, Mediterranean soil, Phosphat Solubilizing Bacteria, *Rhizobium sp.*, *Vigna unguiculata*.

I. INTRODUCTION

In Malang Regency, Mediterranean soil develops from limestone, calcareous sandstone, marl and basalt andesite. The Mediterranean land is found in Karst plains, tectonic plains, tectonic hills, lower volcanic slopes, old volcanic plains and old volcanic hills. Soil cross section is medium to very deep, good drainage, slightly fine to fine texture, slightly acidic to alkaline pH, medium to very high CEC and high to very high base saturation (BS), low to medium in C-organic and total N content., P₂O₅ (medium to high) and total K₂O low to moderate (Yatno *et al.*, 2017). Mediterranean soil has problems with being too alkaline and low in organic matter, macro and micro nutrient levels as well as low water retention capacity and poor drainage (Tri Pamungkas, 2021). Red Mediterranean soil derived from limestone has a higher soil pH value than soil derived from sandstone. The main problem with Mediterranean soil is the high soil pH which will give rise to direct problems, namely suitability for plants and availability of nutrients, namely low availability of microelements and phosphate because they are bound by Ca²⁺ (Supriyadi, 2007). Soil quality and fertility in Mediterranean calcareous agricultural soils are very important to be managed so that soil productivity can be sustainable. The development of Mediterranean land requires pioneer plants that can increase soil fertility, one of which is legumes. Legume crops can be a practical solution for improving calcareous soils because they contribute to nitrogen fixation, organic matter accumulation, and improved soil health. The results of Tri Pamungkas' research showed that the addition of organic fertilizer significantly increased the growth of plant height, number of leaves, and stem diameter of sugar cane seedlings planted in Mediterranean soil. Legume crops are useful as human and animal feed, wood energy, and as soil-improving components of agricultural and agro forestry systems through its association with bio-fertilizers (Havugimana *et al.*, 2016).

Cowpea is included in the minor legume group which has prospects for future development as a source of food, animal feed, green manure, industry, and health. It belongs to the Leguminosae family and as important legume crop growing across the world mainly intropical and subtropical regions (Papa *et al.*, 2020; Kebede and Bekeko, 2020). It is classified as food crops,

feed and industrial raw materials. According to Safitri et al. (2016) stated that cowpea seeds contain a lot of nutrition, every 100 g contains 24.4 g protein, 56.6 g carbohydrates, 1.9 g fat, 481 mg calcium, 399 mg phosphorus, and 2.68 g phytic acid. Cowpea also has a higher vitamin B1 content than green beans. The potential for cowpea seeds is quite high, which can reach 1.5 - 2 tons/ha depending on the variety, location, growing season and cultivation applied. Cowpea has the advantage of having a low fat content so it can minimize the negative effects of fatty food products (Saputro et al., 2015).

Cowpea is a leguminous plant that has also the ability to bind N_2 from the atmosphere. The ability of cowpea to fix nitrogen through biological nitrogen fixation is a cheap and sustainable alternative to inorganic fertilizers. This is because the performance of cowpea depends on the rhizospheric characteristics; hence, it is able to form a beneficial association with microorganisms present in the rhizosphere (Abdel-Fattah et al., 2016). The potency of N_2 fixation by cowpea can be optimized by inoculation of *Rhizobium* sp. Morphologically, cowpea roots spread to a depth of soil between 30-60 cm and can be in symbiosis with *Rhizobium* sp. to bind free Nitrogen (N_2) from the air which then forms root nodules. Inoculation of *Rhizobia* isolates significantly increased the number of root nodules, compared to plants that were not inoculated (Kebede et al., 2020). Symbiosis between cowpea and *Rhizobium* sp. can increase the availability of N to support plant growth and production. *Rhizobium* sp. is one of the bacteria that is very useful for agriculture which acts as a nitrogen fixer from the atmosphere. Nitrogen is one of the main elements in the production of food crops, especially legumes. By fixing nitrogen symbiotically, a cheap source is obtained and can help reduce production costs, especially on infertile soils (Suryantini, 2015). The development of cowpea plants in calcareous areas has several obstacles related to the level of soil fertility. In the calcareous soil, high levels of calcium can also cause problems with the presence of soil potassium. Clay colloids can adsorb and exchange base cations depending on their valence. Cations with three and two valences are more strongly adsorbed by colloids compared to cations with one valence (Putra and Hanum, 2018). As a result, potassium which has a valence of one, will be suppressed by high Ca in calcareous soils which has a valence of two. In calcareous or normal soils, P gets immobilized by cations such as Ca^{2+} to form a complex calcium phosphate ($Ca_3(PO_4)_2$). Nearly, 70 to 90% of phosphorus fertilizers applied to soils is fixed by cations and converted inorganic P (Walpola and Yoon, 2012).

Optimizing the role of cowpeas as a pioneer crop planted in Mediterranean soil requires agronomic practise, one of which is the application of biological fertilizer to help increase the availability of nutrients in the soil, especially elements of N and P to support growth and production. However, the availability of P in the soil is lacking, even though the total amount of P in the soil is high through repeated fertilization applications. This is due to the nature of P which is easily fixed by metals, colloids and calcium which makes it unavailable for plants. Agricultural soils in general also face constraints on the availability of P elements for plants. Even though fertilization has been done many times for each planting, because P is tightly bound by soil colloids, the availability of P element in the soil is low. Likewise with soils that have high Al-dd content, generally also have constraints on the availability of P elements. Low availability of phosphorus (P) in calcareous soils can effect sustainability of improvement in cereals crops yield. A higher amount of calcium in calcareous soils precipitates the P, thus making it immobile in soil. Inoculation of phosphate solubilizing bacteria (PSB) could be helpful in the sustainable management of immobile P in soil (Wahid et al., 2020).

An alternative to increase the efficiency of fertilization in overcoming P fixation by Al, Fe and Ca is by utilizing phosphate solubilizing microorganisms. Phosphate solubilizing bacterial inoculants are microbes given to plants for the process of dissolving P which is bound to become a form available to plants. It play an important role in the biogeochemical cycle of soil P, including dissolution, mineralization, and immobilization of P, thereby increasing P bioavailability and plant nutrition. Root-associated PSB, can modulate plant root systems and stimulate root absorptive capacity and nutrient uptake, including P acquisition (Bargaz et al., 2021). PSB facilitates the conversion of stable phosphorus to active phosphorus and increases the residual phosphorus fraction in compost (Sun et al., 2023). *Bacillus subtilis* is one of PSB which is able to decompose P bonds in the soil and can enhance plant P uptake through remobilization of recalcitrant forms of P in soil. The results of research conducted by Batool and Iqbal (2019) on *Triticum aestivum* L., recommend the use of PSB as biofertilizer, as an alternative to chemical fertilizer.

Providing N and P nutrients in calcareous soil requires technology in the form of the use of biofertilizers which can support the availability of nutrients needed by Cowpea plants to support their growth and production. This study aims to determine the optimal concentration of application of Phosphate solubilizing bacteria and *Rhizobium* sp. to increase biology soil fertility and the growth of cowpea cultivated in Alfisol or Mediterranean soils.

II. MATERIALS AND METHODS

2.1 Study area and soil characteristic

The research was conducted at the Screenhouse of the Faculty of Agriculture, Wisnuwardhana University, Madyopuro Village, Kedungkandang District, Malang City, which is located at an altitude about 500 m above sea level. The type of soil used is the Alfisol or Mediterranean taken from the Bantur area of Malang Regency. The soil used has the following characteristics (primary data from laboratory analysis): pH (H₂O)=7.6 (slightly alkaline), C-Org= 0.73% (very low), N-tot= 0.08% (very low); P-Olsen= 34 ppm (medium); K-dd= 0.24 cmol(+)/kg (low); Ca-dd= 10.20 cmol(+)/kg (high).

2.2 Data collection

Growth parameters observed included plant height, number of root nodules, plant fresh weight, shoot and root dry weight, total dry weight, and weight of 100 seeds, soil PSB population and *Rhizobium sp.* population. Plant height was measured from the base of stem to the top of plant shoot using ruler. The number of root nodules was measured by removing roots at the end of the vegetative phase. The number of effective root nodules is calculated from the pink nodules in color. At the age of 70 days after planting, three samples cowpea plants were taken in each treatment in each replication, then were measured the total fresh weight, dry weight of roots and shoots, and total dry weight of plants. The plant samples were washed then rinsed with distilled water to remove dust and soil on the surface. After the total fresh weight was calculated, then the roots and shoots plant were dried in the oven at 70 °C for 48 hours and weighed to measure the root dry weight and shoot dry weight (Rahayu et al., 2022). The weight of 100 seeds was measured by weighing 100 cowpea seeds taken from each sample plant on each experimental unit after harvest. Calculation of microbial populations of *Rhizobium* used the Selective Medium Total Plate Count (TPC) method. A total of 10 grams of sample was suspended in 90 ml of solution sterile physiological saline, followed by serial dilution using 9 ml of sterile physiological saline to level dilution 10⁻⁵. A total of 100 l of the suspension then spread into Yeast Mannitol (YM) Agar selection medium added with Congo Red, then incubated 3-10 days at room temperature room (28-30 °C). Its growth is observed and its population calculated based on the Most Probable method Number (MPN). Calculation of the population of Phosphate Solubilizing Bacteria using the Selective Medium Total Plate Count (TPC) method. A total of ten grams of sample was suspended in 90 ml of solution sterile physiological saline, then serial dilutions were carried out using 9 ml of sterile physiological saline to a 10⁻⁵ dilution. A total of 100 l of the suspension was then spread over the selection medium Pikovskaya Agar, and incubated for 7 days at room temperature (28 + 2 °C). Growth was observed and the number of colonies counted has a clear zone around the colony based on the MPN method (Peraturan Menteri Pertanian, 2011). Soil chemical properties were analyzed consist of pH (H₂O 1:1), C organic (Walkley and Balck), Nitrogen (Kjeldahl), P (Olsen), K dan Ca (NH₄OAC pH 7.0) (Agricultural Research ad Development Agency, 2012).

2.3 Data analysis

This experiment used a randomized block design, consisting of 9 treatments, namely: control (without PSB and *Rhizobium sp.* inoculants); PSB inoculant concentration of 10 ml/L and *Rhizobium sp.* 10 ml/kg seed; PSB inoculant concentration of 10 ml/L and *Rhizobium sp.* 20 ml/kg seed; PSB inoculant concentration of 10 ml/L and *Rhizobium sp.* 30 ml/kg seed; PSB inoculant concentration of 20 ml/L and *Rhizobium sp.* 10 ml/kg seed; PSB inoculant concentration of 20 ml/L and *Rhizobium sp.* 20 ml/kg seed; PSB inoculant concentration of 20 ml/L and *Rhizobium sp.* 30 ml/kg seed; PSB inoculant concentration of 30 ml/L and *Rhizobium sp.* 10 ml/kg seed; PSB inoculant concentration of 30 ml/L and *Rhizobium sp.* 20 ml/kg seed; PSB inoculant concentration of 30 ml/L and *Rhizobium sp.* 30 ml/kg seed. Each treatment combination was repeated three times. Observational data were analyzed at the 5% level of variance, followed by the 5% BNT test if there was a significant effect of the treatment.

III. RESULTS AND DISCUSSION

3.1 Plant Growth and Production of Cowpea

The results of the varians analysis showed that there was a significant effect on the concentration of Phosphate Solubilizing Bacteria (PSB) and *Rhizobium sp.* on cowpea plant height. Table 1. shows that the height of cowpea plants was significantly different between those treated with phosphate solubilizing bacteria (PSB) and *Rhizobium* bacteria and those without treatment (control). From the first to the ninth week of observation, the average height of cowpea plants in the control showed significantly different and lower than the cowpea applied with PSB and *Rhizobium*. Until the end of the observation, which was 63 days after planting (DAP), the cowpea plants grown on media with the inoculation of PSB 20 ml/l and *Rhizobium* 30 ml/kg, produced the highest growth in plant height (155.33 cm) compared to the other treatments. In fast growth, especially in

the vegetative phase, plants need N elements which play an important role in cell division, the formation of amino acids and proteins, chlorophyll and many potential enzymes that are important for photosynthesis and plant growth (Fitriatin *et al.*, 2017), as well as important for stimulating root growth and increasing the uptake of other nutrients for plants (Nget *et al.*, 2022). So that with higher Rhizobium inoculation it helps the availability of N in the soil.

TABLE 1
PLANT HEIGHT (CM) OF COWPEA (*VIGNA UNGUICULATA*) AS INOCULATION OF PHOSPHATE SOLUBILIZING BACTERIA AND RHIZOBIUM SP.

PSB and Rhizobium Concentration	Plant height (cm)																	
	7 dap		14 dap		21 dap		28 dap		35 dap		42 dap		49 dap		56 dap		63 dap	
0 ml/L and 0 ml/kg	10,17	a	13,17	a	19,17	a	26,50	a	30,83	a	40,83	a	56,67	a	62,67	a	69,33	a
10 ml/L and 10 ml/kg	17,67	b	26,17	b	59,50	c	74,50	b	91,00	b	112,33	bc	106,17	b	110,50	b	116,00	b
10 ml/L and 20 ml/kg	18,17	b	25,33	b	61,50	c	68,17	b	94,50	b	111,83	bc	127,17	c	132,83	bc	136,83	bc
10 ml/L and 30 ml/kg	18,83	b	28,33	b	63,33	c	75,83	b	96,00	b	113,17	bc	120,67	bc	126,83	bc	134,67	bc
20 ml/L and 10 ml/kg	16,67	b	25,33	b	49,83	bc	76,83	b	91,33	b	111,83	bc	133,50	c	136,83	bc	146,17	c
20 ml/L and 20 ml/kg	17,50	b	27,33	b	57,00	bc	79,33	b	96,17	b	92,17	b	131,17	c	136,17	bc	149,50	c
20 ml/L and 30 ml/kg	18,00	b	27,67	b	61,50	c	81,17	b	101,17	b	117,83	c	136,17	c	141,17	c	155,33	c
30 ml/L and 10 ml/kg	18,00	b	24,67	b	39,00	b	75,50	b	101,83	b	126,33	c	134,67	c	138,33	c	144,67	c
30 ml/L and 20 ml/kg	18,50	b	24,83	b	52,67	bc	75,33	b	94,17	b	117,67	c	125,50	c	128,67	bc	142,33	c
30 ml/L and 30 ml/kg	16,50	b	29,00	b	56,83	bc	77,67	b	98,33	b	119,50	c	127,83	c	133,17	bc	143,83	c
BNT 5%	3,005		6,68		18,41		19,13		20,44		22,52		16,45		27,24		25,28	

Note: the numbers followed by the same letter in the same column are not significantly different in the Least Significant Difference Test (LSD) at the 5% level. PSB = Phosphate Solubilizing Bacteria; PSB = Phosphate Solubilizing Bacteria; hst = days after planting; dap = day after planting

Table 2 shows that the control treatment and lower concentration of PSB and Rhizobium, namely 10 ml/l and Rhizobium 10 ml/kg, tended to produce an average weight of 100 cowpea seeds lower than cowpea plants which were inoculated with the higher concentration of PSB (20 ml/l and 30 ml/l) and Rhizobium concentration (20 ml/kg and 30 ml/kg). The results of (Nyaga and Njeru, 2020) research showed that rhizobia inoculation significantly increased cowpea yields where inoculation with native isolates recorded 22.7% increase in yield when compared to uninoculated control in the first season and 28.6% increase in yield in the second season.

TABLE 2
WEIGHT OF 100 SEEDS, NUMBER OF ROOT NODULES, ROOT LENGTH OF COWPEA (*VIGNA UNGUICULATA*) AS INOCULATION OF PHOSPHATE SOLUBILIZING BACTERIA AND RHIZOBIUM SP.

PSB and Rhizobium Concentration	100 seed weight (g)		The Number of root nodule		Root length (cm)	
0 ml/L and 0 ml/kg	7,03	A	1,83	A	9,00	a
10 ml/L and 10 ml/kg	7,17	A	2,17	A	9,33	ab
10 ml/L and 20 ml/kg	7,75	abc	4,00	Ab	12,00	bcde
10 ml/L and 30 ml/kg	8,32	bc	5,00	Bc	12,83	cde
20 ml/L and 10 ml/kg	7,20	Ab	7,00	Cd	10,00	abc
20 ml/L and 20 ml/kg	10,45	D	10,00	E	11,00	abcd
20 ml/L and 30 ml/kg	12,14	E	11,00	Ef	11,00	abcd
30 ml/L and 10 ml/kg	8,65	C	9,00	De	10,83	abcd
30 ml/L and 20 ml/kg	11,77	E	10,00	E	13,00	de
30 ml/L and 30 ml/kg	12,27	E	13,33	F	14,00	e
BNT 5%	1,13		2,34		2,88	

Note: The numbers followed by the same letter in the same column are not significantly different in the Least Significant Difference Test (LSD) at the 5% level. PSB = Phosphate Solubilizing Bacteria.

Observation of the root nodules number showed that the control treatment and the low concentration of PSB and Rhizobium respectively 10 ml/l and 10 ml/kg, tended to produce number of cowpea nodules as 1-2 root nodules. This result showed lower than the higher concentration of PSB (20 ml/l and 30 ml/l) and Rhizobium (20 ml/kg and 30 ml/kg). The highest number of root nodules was produced in the inoculation of PSB 30 ml/l and Rhizobium 30 ml/kg and the application of PSB 20 ml/l by soaking the seeds in Rhizobium 30 ml/kg. The results of this study are in line with the results of Habete and Buraka (2016) research, that showed the presence of rhizobium bacteria inoculation increased the number of nodules per plant which was significantly different for the two varieties of beans. Furthermore, both N fertilizer application and rhizobium inoculation increased 100-seeds weight and seed yield per ha. For better soil health and reduced production costs, it is advisable to use bacteria treated seeds. The root length parameter showed that the cowpea plants that were inoculated with PSB 30 ml/l and Rhizobium sp. 30 ml/kg produced an average root length longer than the other treatments, while the control cowpea plants produced short root lengths. The result of (Chandra and Kumar, 2008) research showed that combined inoculation treatment of *Rhizobium sp.* + Phosphate Solubilizing Bacteria, PSB (*Bacillus megaterium*) + PGPRs LK-786 (*Kurthia sp.*) produced the highest and significantly more number and dry weight of nodules and plant dry weight of Lentil (*Lens culinaris* L.) than *Rhizobium sp.* alone at different intervals. Combined inoculation of chickpea with Rhizobia and Phosphate Solubilizing Bacteria (PSB), *Bacillus sp.*, and *Enterobacter aerogenes* showed a significant enhancement of chickpea nodulation, biomass production, yields and N, P, and protein content in grains as compared to single inoculation or single application of N or P. Formulation of biofertilizers based on tasted strains could be used for chickpea co-inoculation in P-deficient soils (Benjelloun *et al.*, 2021). Rhizobia in association with a legume host have the exceptional ability to form root nodules. PGPR with rhizobial inoculation enhanced nodulation, growth and yield of lentil (*Lens culinaris*) (19.8 nodules/plant, 70.6 mg nodule dry weight/plant, 1 605kg/ ha) as compared to Rhizobium alone (17.8 nodules/plant, 64.3mg/plant and 1 546kg/ha resp.) and non-inoculated control (13.9nodules/plant, 47.7mg/plant and 1 401kg/ha resp.) (Khanna and Sharma, 2011).

TABLE 3

PLANT FRESH WEIGHT, SHOOT DRY WEIGHT, ROOT DRY WEIGHT, AND TOTAL DRY WEIGHT OF COWPEA (*VIGNA UNGUICULATA*) AS INOCULATION OF PHOSPHATE SOLUBILIZING BACTERIA AND *RHIZOBIUM SP.*

PSB and Rhizobium Concentration	Fresh Weight (g)		Root Dry Weight (g)		Shoot Dry Weight (g)		Total Dry Weight (g)	
0 ml/L and 0 ml/kg	65,17	a	0,61	a	18,79	a	19,39	a
10 ml/L and 10 ml/kg	72,00	ab	0,65	a	24,56	b	25,22	b
10 ml/L and 20 ml/kg	73,67	ab	0,47	a	25,31	b	25,78	b
10 ml/L and 30 ml/kg	75,50	ab	0,60	a	25,84	b	26,43	b
20 ml/L and 10 ml/kg	78,67	bc	0,79	a	26,73	b	27,53	b
20 ml/L and 20 ml/kg	80,67	bc	0,53	a	26,78	b	27,30	b
20 ml/L and 30 ml/kg	91,33	c	0,63	a	31,34	c	31,97	c
30 ml/L and 10 ml/kg	71,33	ab	0,44	a	24,54	b	24,97	b
30 ml/L and 20 ml/kg	74,33	ab	0,58	a	25,34	b	26,02	b
30 ml/L and 30 ml/kg	79,83	bc	0,53	a	27,26	bc	27,94	bc
BNT 5%	12,84		0,35		4,37		4,11	

Note: the numbers followed by the same letter in the same column are not significantly different in the Least Significant Difference Test (LSD) at the 5% level. PSB = Phosphate Solubilizing Bacteria.

Table 3 shows that the fresh weight of cowpea was significantly different between those treated with phosphate solubilizing bacteria and Rhizobium bacteria and those without treatment (control). The control showed plant fresh weight (65 g) lower than cowpea inoculated with PSB and Rhizobium. In cowpea plants inoculated with 20 ml/l PSB and *Rhizobium sp.* 10 ml/kg, 20 ml/l PSB and 20 ml/kg *Rhizobium sp.*, 20 ml/l PSB and 30 ml/kg *Rhizobium* resulted in a higher average fresh weight than those inoculated with 10 ml/l PSB at all of *Rhizobium* concentrations level.

The result showed that there were no significant difference on root dry weight between the plants inoculated with PSB and Rhizobium and the control. Meanwhile, there were significant differences on shoot dry weight and total plant dry weight between control and those applied with PSB and *Rhizobium sp.* The highest shoot dry weight and total dry weight were shown in plants that were inoculated with PSB 20 ml/l and Rhizobium 30 ml/kg. The plant growth needs Nitrogen and Phosphorus as a mineral nutrient. Although Phosphorus is abundance in soil, both in organic and inorganic forms, its availability is limited because P mostly occurs in insoluble forms. The average soil P content is about 0.05% (w/w) but only 0.1% of the total P is available to plants due to its low solubility and soil fixation. An adequate supply of phosphorus during the early phases of plant development is important for supporting the development of the reproductive parts of the plant. This is because P plays an important role in increasing root branching and increasing sturdy plants and resistance to disease. Phosphorus accounts for about 0.2 - 0.8% of the plant's dry weight. With the application of phosphate solubilizing microorganisms (PSM) it is the best environmentally friendly means to help the availability of plant P nutrients (Sharma *et al.*, 2013). The combined inoculation of Arbuscular Mycorrhizal Fungi (AMF) and Phosphate Solubilizing Bacteria (PSB) with ground rock phosphate RP had more potential to improve maize-wheat yields and P uptake comparable to those obtained by using expensive phosphatic fertilizers in P deficient calcareous pH soils (Wahid *et al.*, 2020). The research results of Hashem *et al.* (2019) showed that inoculation of plant roots with rhizobia may result in a small number of nodules depending on the root nodulation process which is based on the exchange of signals between the host and bacteria leading to the formation of rhizobia in the host tissue, nodulation and plant growth through increased day element uptake from the soil.

Rhizobium and phosphate solubilizing bacteria are important for plant nutrition. These microbes also have an important role as plant growth-promoting rhizobacteria (PGPR) in plant biofertilizers. Inoculation of Rhizobium and BPF either singly or both with the use of phosphate fertilizer significantly increased root and shoot weight, plant height, seed yield, seed P content, leaf protein content and leaf sugar content. Inoculation of Rhizobium and BPF either singly or both with the addition of phosphate fertilizer improves seed production 30% - 40% better than using only phosphate fertilizer. The synergistic relationship between rhizobium inoculation and BPF which produces growth regulators such as IAA and GA can improve root growth both in length and root weight which in turn can increase water and nutrient uptake which helps improve plant canopy growth (Afzal and Bano, 2008). The role of *Bacillus subtilis* which plays a direct role in P solubilization, it can help the availability of P nutrients in the soil, and can increase plant growth. Among microbial inoculants, the Rhizobium + PSB was found most effective in terms of nodule number (27.66 nodules plant⁻¹), nodule fresh weight (144.90 mg plant⁻¹), nodule dry weight (74.30mg plant⁻¹), shoot dry weight (11.76 g plant⁻¹), and leghemoglobin content (2.29mg g⁻¹ of fresh nodule) and also showed its positive effect in enhancing all the yield attributing parameters, grain and straw yields of Chickpea (Tagore *et al.*, 2013).

3.2 Population of Soils Phosphate Solubilizing Bacteria and Population of Cowpea Roots *Rhizobium sp.*

The results of laboratory analysis showed that inoculation of Phosphate Solubilizing Bacteria (PSB) *Bacillus subtilis* increases the soil PSB population (Table 4). The highest soil PSB population of 2.31×10^7 CFU/g was obtained from the PSB treatment at concentration of 20 ml/L compared to the control i.e population of 1.98×10^6 CFU/g (Table 4). This is in line with the results of Fitriatin *et al.* (2017) that the PSB population in the Rhizosphere of corn plants can be increased by application of PSM biofertilizer (Phosphate Solubilizing Microorganisms) to 2.0×10^{11} CFU/g compared to control (without PSM biofertilizer) as PSB population of 1.2×10^{11} CFU/g. The use of PSB could significantly increase pH, available phosphorus and several kinds of trace elements both in the rhizosphere and non-rhizosphere soil. The PSB secreted small molecular organic acids to dissolve inorganic phosphorus and changed the soil properties, which changed the rhizosphere microbial community indirectly. PSB can produce various organic acids with low molecular weight, which alter the soil pH, solubilizing the phosphate from acid or alkaline soils (Liu *et al.*, 2020).

Alfisols or Mediterranean soils are a group of red soils caused by high iron content and low humus content (Wijanarko *et al.*, 2007). Organic matter is a source of energy for soil macro and micro fauna. Soils with low organic matter content have an effect on low soil microbiological activity and population. Biological fertility (microbiological population) of soil can be increased by addition of organic matter as well as the addition of biological fertilizers. The results showed that increasing the concentration of Rhizobium to 30 ml/kg of seed could increase the soil Rhizobium population up to 1.29×10^7 CFU/g compared to no application of Rhizobium (control) as a population of 2.87×10^6 CFU/g. The increase in microbial population in the soil can be due to the application of bioinoculants which induce better root proliferation and are responsible for high soil enzyme activity resulting in higher biomass production and extended root exudates (Prathima *et al.*, 2022).

TABLE 4
POPULATION OF SOIL PSB AND POPULATION OF COWPEA ROOT RHIZOBIUM SP.

Treatment	Type of Bacteria	Population of Bacteria (CFU/g)
R0 (Control)	<i>Rhizobium sp.</i>	2,87 x 10 ⁶
R1 (10 ml/kg seeds)	<i>Rhizobium sp.</i>	5,60 x 10 ⁶
R2 (20 ml/kg seeds)	<i>Rhizobium sp.</i>	1,01 x 10 ⁷
R3 (30 ml/kg seeds)	<i>Rhizobium sp.</i>	1,29x 10 ⁷
B0 (Control)	PSB <i>Bacillus subtilis</i>	1,98 x 10 ⁶
B1 (10 ml/L)	PSB <i>Bacillus subtilis</i>	3,14 x 10 ⁶
B2 (20 ml/L)	PSB <i>Bacillus subtilis</i>	2,31 x 10 ⁷
B3 (30 ml/L)	PSB <i>Bacillus subtilis</i>	9,61 x 10 ⁶

3.3 Nitrogen and Phosphorus of Soils

Table 5 shows that the Nitrogen total in soils was significantly different between those treated with phosphate solubilizing bacteria and Rhizobium bacteria and those without treatment (control). Planting media and plants inoculated with PSB 10 ml/L and 30 ml/kg *Rhizobium sp.*, PSB 20 ml/L and 30 ml/kg *Rhizobium sp.*, PSB 30 ml/L and 30 ml/kg *Rhizobium sp.*, resulted N total in the soils was higher compared to other treatments. The research result of Wei et al. (2023) showed that long-term co-application of rhizobium and PK (phosphorus and potassium fertilization) promoted soybean nodule dry weight by 33.94% compared with PK + N (nitrogen and PK fertilization), and increased soybean yield by average of 32.25%, 5.90%, and 5.00% compared with CK (non-fertilization control), PK, and PK + N, respectively. The main effect of two rhizobium strain (Faba bean, Fb17 and Fb18) positively improved soil porosity, but soil bulk density was negatively influenced. Soil chemical parameters such as organic carbon, total N, available P, available sulfur, Ca, Mg, and K were positively influenced (Chimdi et al., 2022). Inoculation Rhizobium can increase the availability of N in the soils through nitrogen fixation capability. The research of Oktaviani et al. (2017) showed that combination of Rhizobium and compot 200 g resulted highest value on the total soil N. The highest N soil is produced from combination treatment of 0.245 g, 200 g and Rhizobium compost as much as 25g.

Table 5 also shows that application of 30 ml/L PSB resulted in significantly higher P-total compared to controls and lower PSB concentrations. Total soil P levels in the media and plants applied were 30 ml/L PSB and 20 ml/kg *Rhizobium sp.* showed higher results than the control (without PSB and Rhizobium application) and other treatments. This is very possible because in calcareous soil Ca fixation of soil P becomes a form that is insoluble in the soil solution, and as a consequence is not available to plants. It could be mobilized, converted into soluble P forms using of PSM (Kalayu, 2019). Soil phosphate solubilizing bacteria (PSB) inoculation with mineral P can improved postharvest soil fertility relative to pre-harvest by improving soil organic matter from 0.61% to 0.70%, lowering pH from 7.74 to 7.68, and improving soil total N from 0.04 to 0.09%, ABDTPA-extractable P from 2.07 to 3.44 mg.kg⁻¹, and potassium (K) concentrations from 100.27 to 129.45 mg.kg⁻¹ (Khan et al., 2022).

TABLE 5
THE CONTENT OF NITROGEN AND PHOSPHORUS SOILS AS INOCULATION OF PHOSPHATE SOLUBILIZING BACTERIA AND RHIZOBIUM SP.

PSB <i>Bacillus subtilis</i> and <i>Rhizobium sp.</i> concentration	N-Tot (%)		P-Tot (mg.kg ⁻¹)	
0 ml/L and 0 ml/kg	0,07	a	208	a
10 ml/L and 10 ml/kg	0,21	b	421	b
10 ml/L and 20 ml/kg	0,28	b	454,67	bc
10 ml/L and 30 ml/kg	0,42	c	611,67	bcd
20 ml/L and 10 ml/kg	0,21	b	470,33	bc
20 ml/L and 20 ml/kg	0,28	b	623,33	cd
20 ml/L and 30 ml/kg	0,42	c	616	cd
30 ml/L and 10 ml/kg	0,21	b	604	bcd
30 ml/L and 20 ml/kg	0,28	b	943	e
30 ml/L and 30 ml/kg	0,42	c	723	d
BNT 5%	0,07		192,85	

Note: the numbers followed by the same letter in the same column are not significantly different in the Least Significant Difference Test (LSD) at the 5% level. PSB = Phosphate Solubilizing Bacteria.

IV. CONCLUSION


Inoculation of Phosphate Solubilizing Bacteria (PSB) and *Rhizobium sp.* to cowpea (*Vigna unguiculata*) grown on Alfisol or Mediterranean soils, has an influence on the growth and yield of cowpea plants, namely cowpea plant height, number of root nodules, fresh weight per plant, shoot dry weight and total plant dry weight and average weight of 100 seeds. It's also influence on increasing of soils PSB population and cowpea roots Rhizobium. Total of N and P soils increased by inoculation of PSB and Rhizobium.

The inoculation of PSB 20 ml/l and 30 ml/l and Rhizobium 10 ml/kg, 20 ml/kg and 30 ml/kg resulted in higher plant height, plant fresh weight, shoot dry weight, total dry weight, and weight of 100 seeds of cowpea higher than the other treatments. The highest number of root nodules was produced in the treatment of PSB 30 ml/l and Rhizobium 30 ml/kg. The highest soil PSB population was resulted in the treatment of 20 ml/L and the highest population of cowpea roots *Rhizobium sp.* were produced in the 30 ml/kg treatment. The inoculation of Rhizobium 30 ml/kg produced the highest N-total soil, meanwhile inoculation of PSB 30 ml/L produced the highest P-total soils.

REFERENCES

- [1] Abdel-Fattah, G. M. *et al.* (2016) 'The Impact of the Arbuscular Mycorrhizal Fungi on Growth and Physiological Parameters of Cowpea Plants Grown under Salt Stress Conditions', *International Journal of Applied Sciences and Biotechnology*, 4(3), pp. 372–379. doi: 10.3126/ijasbt.v4i3.15775.
- [2] Afzal, A. and Bano, A. (2008) 'Rhizobium and phosphate solubilizing bacteria improve the yield and phosphorus uptake in wheat (*Triticum aestivum*)', *International Journal of Agriculture and Biology*, 10(1), pp. 85–88.
- [3] Agency, A. R. and D. (2012) *Technical Guidelines for Chemical Analysis of Soil, Plant, Water and Fertilizer*. Ministry of Agriculture of Republic of Indonesia (in Indonesian).
- [4] Bargaz, A. *et al.* (2021) 'Benefits of phosphate solubilizing bacteria on belowground crop performance for improved crop acquisition of phosphorus', *Microbiological Research*. Elsevier GmbH, 252(May), pp. 1–21. doi: 10.1016/j.micres.2021.126842.
- [5] Batool, S. and Iqbal, A. (2019) 'Phosphate solubilizing rhizobacteria as alternative of chemical fertilizer for growth and yield of *Triticum aestivum* (Var. Galaxy 2013)', *Saudi Journal of Biological Sciences*. King Saud University, 26(7), pp. 1400–1410. doi: 10.1016/j.sjbs.2018.05.024.
- [6] Benjelloun, I. *et al.* (2021) 'Co-inoculation of mesorhizobium ciceri with either bacillus sp. Or enterobacter aerogenes on chickpea improves growth and productivity in phosphate-deficient soils in dry areas of a mediterranean region', *Plants*, 10(3), pp. 1–15. doi: 10.3390/plants10030571.
- [7] Chandra, R. and Kumar, R. (2008) 'Influence of PGPR and PSB on Rhizobium leguminosarum Bv. viciae Strain Competition and Symbiotic Performance in Lentil', *World Journal of Agricultural Sciences*, 4(3), pp. 297–301.
- [8] Chimdi, A., Negasa, D. and Chala, G. (2022) 'Effects of Rhizobium Inoculation and P Fertilizer Levels on Selected Soil Properties, Yield, and Yield Components of Faba Bean (*Vicia faba* L.): The Case of Abuna Gindeberat, West Shewa Zone, Ethiopia', *Applied and Environmental Soil Science*, 2022, pp. 1–10. doi: 10.1155/2022/3635989.
- [9] Fitriatin, B. N., Agustina, M. and Hindersah, R. (2017) 'Populasi Bakteri Pelarut Fosfat , P-Potensial Dan Hasil Jagung Yang Total Phosphate Solubilizing Bacteria (Psb), Soil Potential P And Yield Of Maize (*Zea mays* . L) Affected By The MPF Application Grown On Jatinangor Ultisols', *Jurnal Agrologia*, 6(2), pp. 75–83.
- [10] Habete, A. and Buraka, T. (2016) 'Effect of Rhizobium Inoculation and Nitrogen Fertilization on Nodulation and Yield Response of Common Bean (*Phaseolus vulgaris* L.) at Boloso Sore, Southern Ethiopia', *Journal of Biology, Agriculture and Healthcare*, 6(13), pp. 72–75. Available at: www.iiste.org.
- [11] Hashem, A., Tabassum, B. and Fathi Abd_Allah, E. (2019) 'Bacillus subtilis: A plant-growth promoting rhizobacterium that also impacts biotic stress', *Saudi Journal of Biological Sciences*. King Saud University, 26(6), pp. 1291–1297. doi: 10.1016/j.sjbs.2019.05.004.
- [12] Havugimana, E. *et al.* (2016) 'Role of dual inoculation of Rhizobium and arbuscular mycorrhizal (AM) fungi on pulse crops production', *Walailak Journal of Science and Technology*, 13(1), pp. 1–7.
- [13] Kalayu, G. (2019) 'Phosphate Solubilizing Microorganisms: Promising Approach as Biofertilizers', *International Journal of Agronomy*, 2019, pp. 1–7.
- [14] Kebede, E. *et al.* (2020) 'Symbiotic effectiveness of cowpea (*Vigna unguiculata* (L.) Walp.) nodulating rhizobia isolated from soils of major cowpea producing areas in Ethiopia', *Cogent Food and Agriculture*. Cogent, 6(1), pp. 1–16. doi: 10.1080/23311932.2020.1763648.
- [15] Kebede, E. and Bekeko, Z. (2020) 'Expounding the production and importance of cowpea (*Vigna unguiculata* (L.) Walp.) in Ethiopia', *Cogent Food and Agriculture*. Cogent, 6(1). doi: 10.1080/23311932.2020.1769805.
- [16] Khan, H. *et al.* (2022) 'Coupling phosphate-solubilizing bacteria (PSB) with inorganic phosphorus fertilizer improves mungbean (*Vigna radiata*) phosphorus acquisition, nitrogen fixation, and yield in alkaline-calcareous soil', *Heliyon*. The Author(s), 8(3), pp. 1–10. doi: 10.1016/j.heliyon.2022.e09081.
- [17] Khanna, V. and Sharma, P. (2011) 'Potential for enhancing lentil (*Lens culinaris*) productivity by co-inoculation with PSB, plant growth-promoting rhizobacteria and Rhizobium', *Indian Journal of Agricultural Sciences*, 81(10), pp. 932–934.

- [18] Liu, J. *et al.* (2020) 'Exogenous phosphorus-solubilizing bacteria changed the rhizosphere microbial community indirectly', *3 Biotech.* Springer International Publishing, 10(4), pp. 1–11. doi: 10.1007/s13205-020-2099-4.
- [19] Nget, R. *et al.* (2022) 'Responses of Soybean Genotypes to Different Nitrogen and Phosphorus Sources: Impacts on Yield Components, Seed Yield, and Seed Protein', *Plants*, 11(3). doi: 10.3390/plants11030298.
- [20] Nyaga, J. W. and Njeru, E. M. (2020) 'Potential of Native Rhizobia to Improve Cowpea Growth and Production in Semiarid Regions of Kenya', *Frontiers in Agronomy*, 2(December), pp. 1–13. doi: 10.3389/fagro.2020.606293.
- [21] Oktaviani, E. and Susanto, D. (2017) 'Effect Of Inoculant Rhizobium And Compost As Amelioration On Growth Of Peanut (*Arachis Hypogaea*) In Soil Post-Coal Mining', *International Journal of Scientific & Technology Research*, 6(02), pp. 4–8.
- [22] Papa, M., Hetharie, H. and Polnaya, F. (2020) 'Keragaan Pertumbuhan dan Produksi Aksesori-Aksesori Kacang Tunggak Lokal Asal Maluku', *Jurnal Budidaya Pertanian*, 16(1), pp. 88–94. doi: 10.30598/jbdp.2020.16.1.88.
- [23] Peraturan Menteri Pertanian (2011) 'Pupuk Organik, Pupuk Hayati Dan Pembenah Tanah', *Peraturan Menteri Pertanian Nomor 70/Permentan/SR.140/10/2011*, pp. 1–109.
- [24] Prathima, M. N. *et al.* (2022) 'Influence of Bioinoculants on Growth and Yield of Cowpea (*Vigna unguiculata* L.) under Field Condition', *Agricultural Research Communication Centre (ARCC) Journals: Legume Research: An International Journal*, 1, pp. 1–6. doi: 10.9734/ijec/2022/v12i1030773.
- [25] Putra, I. A. and Hanum, H. (2018) 'Kajian Antagonisme Hara K, Ca Dan Mg pada Tanah Inceptisol yang Diaplikasi Pupuk Kandang, Dolomit dan Pupuk KCl terhadap Pertumbuhan Jagung Manis (*Zea mays saccharata* L.) Antagonism Study K, Ca And Mg Nutrients On The Inceptisol Soil that Applied Man', *Elkawanie: Journal of Islamic Science and Technology*, 4(1), pp. 23–44.
- [26] Rahayu, Y. S., Maghfoer, M. D. and Wardiyati, T. (2022) 'The potential of intercropping of *Crotalaria juncea* on the reduction of Pb accumulation in *Brassica rapa* and *Phaseolus vulgaris* grown on Pb-contaminated soil', *Journal of Degraded and Mining Lands Management*, 9(4), pp. 3745–3752. doi: 10.15243/jdmlm.2022.094.3745.
- [27] Safitri, F. M. *et al.* (2016) 'Pengembangan getuk kacang tolo sebagai makanan selingan alternatif kaya serat', *Jurnal Gizi dan Dietetik Indonesia (Indonesian Journal of Nutrition and Dietetics)*, 4(2), p. 71. doi: 10.21927/ijnd.2016.4(2).71-80.
- [28] Saputro, D. H., Andriani, M. and Siswanti (2015) 'Karakteristik Sifat Fisik Dan Kimia Formulasi Tepung Kecambah Kacang-Kacangan Sebagai Bahan Minuman', *Jurnal Teknosains Pangan*, 4(1), p. 11. Available at: www.ilmupangan.fp.uns.ac.id.
- [29] Sharma, S. B. *et al.* (2013) 'Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils', *Springer Plus*, 2(587), pp. 1–14. doi: 10.1128/jcm.35.12.3305-3307.1997.
- [30] Sun, L. *et al.* (2023) 'Effects of phosphate-solubilizing bacteria on phosphorus components, humus and bacterial community metabolism during spent mushroom substrate composting', *Environmental Technology and Innovation*. Elsevier B.V., 32, pp. 1–13. doi: 10.1016/j.eti.2023.103341.
- [31] Supriyadi, S. (2007) 'Kesuburan Tanah Di Lahan Kering Madura', *Embryo*, 4(2), pp. 124–131.
- [32] Suryantini (2015) 'Pembintilan dan Penambatan Nitrogen', *Monograf Balitkabi*, pp. 234–250.
- [33] Tagore, G. S. *et al.* (2013) 'Effect of Rhizobium and Phosphate Solubilizing Bacterial Inoculants on Symbiotic Traits, Nodule Leghemoglobin, and Yield of Chickpea Genotypes', *International Journal of Agronomy*, 2013, pp. 1–8. doi: 10.1155/2013/581627.
- [34] Tri Pamungkas, S. S. (2021) 'Pemanfaatan Tanah Mediteranian Sebagai Media Pembibitan Budet Tebu (*Saccharum officinarum* L.) Varietas Bululawang dengan Penambahan Pupuk Kandang pada Dosis yang Berbeda', *Mediagro*, 17(2), pp. 107–119. doi: 10.31942/md.v17i2.4918.
- [35] Wahid, F. *et al.* (2020) 'Sustainable management with mycorrhizae and phosphate solubilizing bacteria for enhanced phosphorus uptake in calcareous soils', *Agriculture (Switzerland)*, 10(8), pp. 1–14. doi: 10.3390/agriculture10080334.
- [36] Walpola, B. C. and Yoon, M. (2012) 'Prospectus of phosphate solubilizing microorganisms and phosphorus availability in agricultural soils: A review', *African Journal of Microbiology Research*, 6(37), pp. 6600–6605. doi: 10.5897/AJMR12.889.
- [37] Wei, W. *et al.* (2023) 'Long-term fertilization coupled with rhizobium inoculation promotes soybean yield and alters soil bacterial community composition', *Frontiers in Microbiology*, 14(May), pp. 1–16. doi: 10.3389/fmicb.2023.1161983.
- [38] Wijanarko, A., Sudaryono and Sutarno (2007) 'Characteristics of Chemical and Physical Soil Alfisols in East Java and Central Java', *Iptek Tanaman Pangan*, 2(2), pp. 214–226.
- [39] Yatno, E. *et al.* (2017) *Atlas Peta Tanah Semi Detail Skala 1 : 50.000 Kabupaten Malang Provinsi Jawa Timur*. 2017th edn. Edited by Sukarman and M. Anda. Malang: Balai Besar Penelitian dan Pengembangan Sumberdaya Lahan Pertanian- Badan Penelitian dan Pengembangan Pertanian- Kementerian Pertanian.



AD Publications

Sector-3, MP Nagar, Rajasthan, India

www.adpublications.org, www.ijoeear.com, info@ijoeear.com